

40 Rec'd PCT/PTO 25 MAR 1998

FORM PTO-1390

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ISIS-2297

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

254-65
U.S. APPLICATION NO. (if known see 37 C.F.R. 1.5)
08/973,381 #2

INTERNATIONAL APPLICATION NO.
PCT/US96/09604

INTERNATIONAL FILING DATE
JUNE 7, 1996

PRIORITY DATE CLAIMED
JUNE 7, 1995

TITLE OF INVENTION COMBINATORIAL LIBRARIES HAVING AMINODIOL MONOMER SUBUNITS

APPLICANT(S) FOR DO/EO/US Normand HEBERT

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☐ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☒ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☐ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) 35 U.S.C. 371(c)(4).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:


11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Associate Power of Attorney

EXPRESS MAIL Mailing Label No. EL066384310US


Date of Deposit: MARCH 25, 1998

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

MAILER BOB INFORZATO

SIGNATURE 

03/27/1998 PVDLPE 00000080 08973381
01 FC:254 65.00 OP

U.S. APPLICATION NO. (if known 37 C.F.R. 1.5) 08/973,381		INTERNATIONAL APPLICATION NO. PCT/US96/09604		ATTORNEY DOCKET NUMBER ISIS-2297	
17. <input checked="" type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Search Report has been prepared by the EPO or JPO.....\$930.00 International preliminary examination fee paid to USPTO (37 CFR 1.482)\$720.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)).....\$790.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$1,070.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4).....\$98.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>				<u>CALCULATIONS</u> <u>PTO USE ONLY</u>	
Surcharge of \$130.00 for furnishing the oath or declaration later than <u> 20 </u> <input checked="" type="checkbox"/> <u> 30 </u> months from the earliest claimed priority date (37 CFR 1.492(e)).				\$130	
Claims	Number Filed	Number Extra	Rate		
Total claims	- 20 =		X \$22.00	\$	
Independent Claims	- 3 =		x \$82.00	\$	
Multiple dependent claims(s) (if applicable)			+ \$270.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$130	
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28). Small Entity Statement was filed on December 3, 1997				\$65	
SUBTOTAL =				\$65	
Processing fee of \$130.00 for furnishing the English translation later than <u> 20 </u> <u> 30 </u> months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$65	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				+	
TOTAL FEES ENCLOSED =				\$65	
				Amount to be: refunded	\$
				charged	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>65</u> to cover the above fee is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. 23-3050 in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-3050. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Michael P. Straher Woodcock Washburn Kurtz Mackiewicz & Norris LLP One Liberty Place - 46th Floor Philadelphia, PA 19103 (215) 568-3100			<div style="text-align: center;">  _____ SIGNATURE Michael P. Straher _____ NAME 38,325 _____ REGISTRATION NUMBER </div>		

FORM PTO-1390

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY DOCKET NUMBER **ISIS-2297**

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DQ/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

U.S. APPLICATION NO. (if known see 37 C.F.R. 1.5)

08/973381INTERNATIONAL APPLICATION NO.
PCT/US96/09604INTERNATIONAL FILING DATE
07 JUNE 1996PRIORITY DATE CLAIMED
07 JUNE 1995TITLE OF INVENTION **COMBINATORIAL LIBRARIES HAVING AMINODIOL MONOMER SUBUNITS**APPLICANT(S) FOR DO/EO/US **HEBERT, Normand**

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) 35 U.S.C. 371(c)(4).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Verified Statement Claiming Small Entity Status

EXPRESS MAIL Mailing Label No. EM155912058US

Date of Deposit: **DECEMBER 3, 1997** *B.A.*

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

MAILER **BOB INFORZATO**SIGNATURE *Bob Inforzato*

U.S. APPLICATION NO. (if known 37 C.F.R. 1.5)	INTERNATIONAL APPLICATION NO. PCT/US96/09604	ATTORNEY DOCKET NUMBER ISIS-2297
---	---	-------------------------------------

<p>17. <u>X</u> The following fees are submitted:</p> <p style="margin-left: 20px;">Basic National Fee (37 CFR 1.492(a)(1)-(5)):</p> <p style="margin-left: 20px;">Search Report has been prepared by the EPO or JPO.....\$930.00</p> <p style="margin-left: 20px;">International preliminary examination fee paid to USPTO (37 CFR 1.482)\$720.00</p> <p style="margin-left: 20px;">No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)).....\$790.00</p> <p style="margin-left: 20px;">Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$1,070.00</p> <p style="margin-left: 20px;">International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4).....\$98.00</p>	<p><u>CALCULATIONS</u> <u>PTO USE ONLY</u></p>
--	---

ENTER APPROPRIATE BASIC FEE AMOUNT =	\$720.00	
--------------------------------------	----------	--

Surcharge of \$130.00 for furnishing the oath or declaration later than <u> 20 </u> <u> 30 </u> months from the earliest claimed priority date (37 CFR 1.492(e)).	\$	
---	----	--

Claims	Number Filed	Number Extra	Rate		
Total claims	26 - 20 =	6	X \$22.00	\$132	
Independent Claims	4 - 3 =	1	x \$82.00	\$82	
Multiple dependent claims(s) (if applicable)			+ \$270.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$934	
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).				\$467	
SUBTOTAL =				\$467	
Processing fee of \$130.00 for furnishing the English translation later than <u> 20 </u> <u> 30 </u> months from the earliest claimed priority date (37 CFR 1.492(f)).				+	\$
TOTAL NATIONAL FEE =				\$467	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				+	
TOTAL FEES ENCLOSED =				\$467	


	Amount to be:	
	refunded	\$
	charged	\$

a. X A check in the amount of \$ 467.00 to cover the above fee is enclosed.

b. Please charge my Deposit Account No. 23-3050 in the amount of \$ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. X The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-3050. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

<p>SEND ALL CORRESPONDENCE TO:</p> <p>STRAHER, Michael P.</p> <p>Woodcock Washburn Kurtz</p> <p>Mackiewicz & Norris LLP</p> <p>One Liberty Place - 46th Floor</p> <p>Philadelphia, PA 19103</p> <p>(215) 568-3100</p>	<p style="text-align: center;"></p> <p style="text-align: center;">SIGNATURE</p> <hr style="border: 0; border-top: 1px solid black; margin: 5px 0;"/> <p style="text-align: center;">STRAHER, Michael P.</p> <p style="text-align: center;">NAME</p> <hr style="border: 0; border-top: 1px solid black; margin: 5px 0;"/> <p style="text-align: center;">38,325</p> <p style="text-align: center;">REGISTRATION NUMBER</p>
---	--

08/973381

COMBINATORIAL LIBRARIES HAVING AMINODIOL MONOMER SUBUNITS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of United States Application Serial No. 08/483,311, filed
5 6/7/95;, which is a continuation-in-part of PCT application Serial No. PCT/US95 /00356 filed 1/11/95, which is a continuation-in-part of United States Application Serial No. 08/180,134, filed 1/11/94; which is a continuation in part
10 of United States Application 08/179,970, filed 1/11/94 which issued on 5/21/96 as U.S. Patent Serial No. 5519134. Each of these patent applications are assigned to the assignee of this application and are incorporated by reference herein.

BACKGROUND OF THE INVENTION

Traditional processes of drug discovery involve
15 the screening of complex fermentation broths and plant extracts for a desired biological activity or the chemical synthesis of many new compounds for evaluation as potential drugs. The advantage of screening mixtures from biological sources is that a large number of compounds are screened
20 simultaneously, in some cases leading to the discovery of novel and complex natural products with activity that could not have been predicted otherwise. The disadvantages are that many different samples must be screened and numerous

- 2 -

purifications must be carried out to identify the active component, often present only in trace amounts. On the other hand, laboratory syntheses give unambiguous products, but the preparation of each new structure requires significant amounts of resources. Generally, the *de novo* design of active compounds based on the high resolution structures of enzymes has not been successful.

It is thus now widely appreciated that combinatorial libraries are useful *per se* and that such libraries and compounds comprising them have great commercial importance. Indeed, a branch of chemistry has developed to exploit the many commercial aspects of combinatorial libraries.

In order to maximize the advantages of each classical approach, new strategies for combinatorial deconvolution have been developed independently by several groups. Selection techniques have been used with libraries of peptides (Geysen, H. M., Rodda, S. J., Mason, T. J., Tribbick, G. and Schoofs, P. G., *J. Immun. Meth.* **1987**, *102*, 259-274; Houghten, R. A., Pinilla, C., Blondelle, S. E., Appel, J. R., Dooley, C. T. and Cuervo, J. H., *Nature*, **1991**, *354*, 84-86; Owens, R. A., Gesellchen, P. D., Houchins, B. J. and DiMarchi, R. D., *Biochem. Biophys. Res. Commun.*, **1991**, *181*, 402-408; Doyle, M.V., PCT WO 94/28424; Brennan, T.M., PCT WO 94/27719); nucleic acids (Wyatt, J. R., et al., *Proc. Natl. Acad. Sci. USA*, **1994**, *91*, 1356-1360; Ecker, D. J., Vickers, T. A., Hanecak, R., Driver, V. and Anderson, K., *Nucleic Acids Res.*, **1993**, *21*, 1853-1856); nonpeptides and small molecules (Simon, R. J., et al., *Proc. Natl. Acad. Sci. USA*, **1992**, *89*, 9367-9371; Zuckermann, R. N., et al., *J. Amer. Chem. Soc.*, **1992**, *114*, 10646-10647; Bartlett, Santi, Simon, PCT WO91/19735; Ohlmeyer, M. H., et al., *Proc. Natl. Acad. Sci. USA*, **1993**, *90*, 10922-10926; DeWitt, S. H., Kiely, J. S., Stankovic, C. J., Schroeder, M.C. Reynolds

- 3 -

Cody, D.M. and Pavia, M.R., *Proc. Natl. Acad. Sci. USA*,
1993, 90, 6909-6913; Cody et al., United States patent
5,324,483; Houghten et al., PCT WO 94/26775; Ellman, United
States patent 5,288,514; Still et al., PCT WO 94/08051;
5 Kauffman et al., PCT WO 94/24314; Carell, T., Wintner, D.
A., Bashir-Hashemi, A. and Rebek, J., *Angew. Chem. Int. Ed.*
Engel., 1994, 33, 2059-2061; Carell, T., Wintner, D. A. and
Rebek, J., *Angew. Chem. Int. Ed. Engel.*, 1994, 33, 2061-
2064; Lebl, et al., PCT WO 94/28028). A review of the above
10 references reveals that the most advanced of these
techniques are those for selection of peptides and nucleic
acids. Several groups are working on selection of
heterocycles such as benzodiazepines. With the exception of
Rebek et al., scant attention has been given to
15 combinatorial discovery of other types of molecules.

The majority of the techniques reported to date
involve iterative synthesis and screening of increasingly
simplified subsets of oligomers. Monomers or sub-monomers
that have been utilized include amino acids, amino acid-like
20 molecules, i.e. carbamate precursors, and nucleotides, both
of which are bifunctional. Utilizing these techniques,
libraries have been assayed for activity in either cell-
based assays, or for binding and/or inhibition of purified
protein targets.

25 A technique, called SURF™ (Synthetic Unrandomization of
Randomized Fragments), involves the synthesis of subsets of
oligomers containing a known residue at one fixed position
and equimolar mixtures of residues at all other positions.
For a library of oligomers four residues long containing
30 three monomers (A, B, C), three subsets each containing 27
compounds would be synthesized (NNAN, NNBN, NNCN, where N
represents equal incorporation of each of the three
monomers). Each subset is then screened in a functional
assay and the best subset is identified (e.g. NNAN). A

- 4 -

second set of subsets is synthesized and screened, each containing the fixed residue from the previous round, and a second fixed residue (e.g. ANAN, BNAN, CNAN, each containing 9 molecules). Through successive rounds of screening and synthesis, a unique sequence with activity in the functional assay can be identified. The SURF™ technique is described in Ecker, D.J., Vickers, T. A., Hanecak, R., Driver, V. & Anderson, K., *Nucleic Acids Res.*, 1993, 21, 1853-1856. The SURF™ method is further described in PCT patent application WO 93/04204, the entire disclosure of which is herein incorporated by reference.

The combinatorial chemical approach that has been most utilized to date, utilizes an oligomerization from a solid support using monomeric units and a defined connecting chemistry, i.e. a solid support monomer approach. This approach has been utilized in the synthesis of libraries of peptides, peptoids, carbamates and vinylogous peptides connected by amide or carbamate linkages or nucleic acids connected by phosphate linkages as exemplified by the citations in previous paragraphs above. A mixture of oligomers (pool or library) is obtained from the addition of a mixture of activated monomers during the coupling step or from the coupling of individual monomers with a portion of the support (bead splitting) followed by remixing of the support and subsequent splitting for the next coupling. In this monomeric approach, each monomeric unit would carry a tethered letter, i.e., a functional group for interaction with the target. Further coupling chemistry that allows for the insertion of a tethered letter at a chemically activated intermediate stage is referred to as the sub-monomer approach.

The diversity of the oligomeric pool is represented by the inherent physical properties of each monomer, the number of different monomers mixed at each coupling, the physical properties of the chemical bonds

- 5 -


arising from the coupling chemistry (the backbone), the number of couplings (length of oligomer), and the interactions of the backbone and monomer chemistries. Taken together, these interactions provide a unique conformation for each individual molecule.

There remains a need in the art for molecules which have fixed preorganized geometry that matches that of targets such as proteins and enzymes, nucleic acids, lipids and other targets. The backbone of such molecules should be rigid with some flexibility, and such molecules should be easy to construct via automated synthesis on solid support.

SUMMARY OF THE INVENTION

In accordance with this invention there are provided oligomeric compounds and libraries of such compounds comprising a plurality of aminodiol monomer subunits joined by linking groups, wherein each of said aminodiol monomer subunits has one of the structures I, II, III, IV, V, VI, VII, VIII, IX, X, or XI;

- 6 -



wherein:


each x is, independently, 0 to 5;

na, nb and nc are each, independently, 0 to 2,

where the sum of na, nb and nc is from 1 to 5;

5 R_1 is -T-L or a base labile protecting group;

T is a single bond, a methylene group or a group having formula:



where:

R_{10} is =O, =S, or =NR₁₁;

R_5 and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;

5 each R_6 , R_7 , R_8 , R_9 , R_{11} , R_{12} and R_{13} are, independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

10 m and n, independently, are 0 to 5;

p is 0 or 1;

q is 1 to about 10;


L is H, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted C₄-C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇ carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and

- 8 -

alkynyl groups;

R_2 is hydrogen or C_1 - C_{10} alkyl;

R_3 and R_4 are independently hydrogen, an acid labile hydroxyl protecting group, a linking group or a conjugate group, wherein said linking group has the formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_0)T_0$;

Y_0 is H or $[Q_2]_j-Z_2$;

T_0 is $[Q_1]_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_4 - C_7 carbocyclo alkyl, C_4 - C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7 - C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H, C_1 - C_2 alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, C_6 - C_{14} aryl, C_7 - C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group; and

provided that at least one of said aminodiol monomer subunits in said oligomeric compound does not have structure III.


Further in accordance with this invention there are provided processes for preparing oligomeric

- 9 -

compounds and libraries of such compounds comprising:

(a) selecting an aminodiol monomer subunit having the structure I, II, III, IV, V, VI, VII, VIII, IX, X, or XI:

- 10 -



wherein:

each x is, independently, 0 to 5;

na , nb and nc are each, independently, 0 to 2, where the sum of na , nb and nc is from 1 to 5;

5 R_1 is a base labile protecting group;

R_2 is hydrogen or C_1 - C_{10} alkyl; and

one of R_3 or R_4 is hydrogen or an activated phosphite group and the other of R_3 or R_4 is an acid labile hydroxyl protecting group;

10 (b) attaching said aminodiol monomer subunit to a

- 11 -


solid support to form a solid support bound aminodiol monomer subunit;

(c) treating said acid labile hydroxyl protecting group with a dilute acid to form a free hydroxyl group,

5 (d) reacting said free hydroxyl group with a further aminodiol monomer subunit having structure I, II, III, IV, V, VI, VII, VIII, IX, X, or XI, thereby forming an oligomeric compound bound to said solid support, said oligomeric compound containing a phosphite linkage;

10 (e) optionally iteratively repeating steps (c) and (d) to increase the length of the oligomeric compound bound to said solid support;

(f) optionally, prior to step (c) or after step (d) oxidizing said phosphite linkage to form a phosphate linking group wherein said linking groups are selected
15 having formula:



wherein:

- 20 J_1 is =O or =S;
 J_2 is OH or N(Y₀)T₀;
 Y_0 is H or [Q₂]_j-Z₂;
 T_0 is [Q₁]_k-Z₁, or together Y₀ and T₀ are joined in a nitrogen heterocycle;
 Q_1 and Q_2 independently, are C₂-C₁₀ alkyl, C₂-C₁₀
 25 alkenyl, C₂-C₁₀ alkynyl, C₄-C₇ carbocyclo alkyl or alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C₇-C₁₄ aralkyl;
 j and k independently, are 0 or 1;
 Z_1 and Z_2 independently are H, C₁-C₂ alkyl, C₂-C₂₀
 30 alkenyl, C₂-C₂₀ alkynyl, C₆-C₁₄ aryl, C₇-C₁₅ aralkyl, halogen,


- 12 -

CH=O, OR₁₂, SR₁₂, NR₁₂R₁₃, C(=NH)NR₁₂R₁₃, CH(NR₁₂R₁₃),
 NHC(=NH)NR₁₂R₁₃, CH(NH₂)C(=O)OH, C(=O)NR₁₂R₁₃, C(=O)OR₁₂, a metal
 coordination group, a reporter group, a nitrogen-containing
 heterocycle, a purine, a pyrimidine, a phosphate group, a
 5 polyether group, or a polyethylene glycol group; and

(g) prior to step (e) or after step (f) contacting
 said solid support bound aminodiol monomer subunit or said
 support bound oligomeric compound with a base to remove said
 base labile amino protecting group to form the solid support
 10 bound aminodiol monomer subunit or support bound oligomeric
 compound having a free amine, and derivatizing said free
 amine with a group of the formula -T-L;

wherein:

T is a single bond, a methylene group or a group having
 15 formula:



where:

R₁₀ is =O, =S, or =NR₁₁;
 20 R₅ and E, independently, are a single bond, CH=CH,
 C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;

each R₆, R₇, R₈, R₉, R₁₁, R₁₂ and R₁₃ are,
 independently, H, alkyl or haloalkyl having 1 to about 10
 carbon atoms, alkenyl having 2 to about 10 carbon atoms,
 25 alkynyl having 2 to about 10 carbon atoms, or aryl having 7
 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

p is 0 or 1;

q is 1 to about 10;

30 L is H, substituted or unsubstituted C₂-C₁₀ alkyl,
 substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or
 unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted
 C₄-C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇

- 13 -

carbocyclic alkenyl, substituted or unsubstituted C₄-C₇,
carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄, aryl,
an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or
sulfur atoms, a nitrogen containing heterocycle, a sulfur
5 containing heterocycle, an oxygen containing heterocycle, a
metal coordination group, a conjugate group, halogen,
hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH),
amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine
(NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O),
10 nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl
(CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl,
N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂),
azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide
(SO), sulfone (SO₂), sulfide (S-), disulfide (S-S), silyl, a
15 nucleosidic base, an amino acid side chain, a carbohydrate,
a biopharmaceutically active moiety, or group capable of
hydrogen bonding where the substituent groups are selected
from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl,
nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and
20 alkynyl groups;

(h) optionally repeating steps (c) and (d)
followed by step (g) to increase the length of the
oligomeric compound bound to said solid support;

(i) treating said oligomeric compound bound to
25 said solid support with acid to deprotect any protecting
groups; and

(j) cleaving said oligomeric compound from said
solid support.

BRIEF DESCRIPTION OF THE FIGURES

30 Figure 1A and Figure 1B are flow schemes illustrating
certain processes of the inventions.

DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention provides combinatorial libraries of phosphate linked aminodiol monomer subunits having phosphodiester, phosphorothioate, and phosphoramidate linkages. As used in the context of the present invention an aminodiol monomer subunit (monomer subunit or backbone segment) is a cyclic or acyclic compound having a protected reactive amino group either primary or secondary and two hydroxyls that can be either primary or secondary. One of the hydroxyls is protected with an acid labile protecting group and the other is in the form of the free hydroxyl or in the form of an H-phosphonate. The monomer subunits of the invention are coupled using H-phosphonate chemistry to form oligomeric compounds which are substituted with diverse functional groups.

Oligomeric compounds of the invention have reactive sites, also referred to as combinatorial sites, that may be combinatorialized with diverse functional groups. Sites that are available for combinatorializing include reactive amino groups, terminal hydroxyl groups, and H-phosphonate phosphorous groups that can be oxidized with, for example, a) carbon tetrachloride and primary or secondary amines to form a phosphoramidate, b) iodine and water to form phosphodiester, or c) sulfur in carbon disulfide to form phosphorothioates and terminal hydroxyls. Terminal hydroxyls may also be substituted with a variety of functional groups such as nucleotides or nucleosides to give a chimeric compound or may include any of a variety of other groups such as conjugates, and reporter groups.

Functional groups may be attached directly to combinatorial sites or may include a tethering group to alter their orientation in space. The functional groups are attached to the backbone segment and phosphoramidate moiety with or without intervening tethering groups. Tethering groups, as used in the context of this invention, are bivalent or polyvalent groups that have a primary or secondary amine or

- 15 -

other suitable group to react with an H phosphonate backbone segment of the invention together with a second functional group capable of binding a "letter". Such tethers can be used to position "letters" in space with respect to the linear backbone of the oligomeric compound synthesized or to link letters that themselves do not include an amine group - necessary to form a phosphoramidate linkage - as an inherent part of the letter. A particularly preferred group of compounds, when substituted with an appropriate amine functional group where necessary, useful as tethering groups include, but are not limited to C₂-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₄-C₇ carbocyclo-alkyl or alkenyl, heterocycles, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, polyalkylene glycols and C₇-C₁₄ aralkyl groups. Other representative tethers useful in the practice of the invention are disclosed in United States application serial number 08/116,801, filed Sept. 3, 1993, entitled "Thiol-Derivatized Nucleosides and Oligonucleosides" and United States application serial number 117,363, filed Sept. 3, 1993, entitled "Amine-Derivatized Nucleosides and Oligonucleosides", the disclosures of which are hereby incorporated by reference.

The present invention provides for the addition of functional groups onto the backbone of the oligomeric compounds on the solid support. In contrast to previous methods, the synthesis of a large set of monomer subunits bearing various functional groups is not necessary. In one aspect the oligomeric compounds of the invention are composed of three components. The preparation of the combinatorial libraries begins with an aminodiol monomer subunit attached to the solid support directly through a linker stable to the synthesis conditions, but cleavable to release the compound into solution at the end of the synthesis. Preferred linkers include esters, particularly succinic acid. Alternatively, the monomers can be coupled

- 16 -

to a constant moiety attached to the CPG, such as DMT ethylene glycol or a similar diol. Other attachment points to the solid support are possible, for example: nucleosides, amino acids or other groups imparting pharmacokinetic, pharmacodynamic or other desirable properties.

The backbone segments are a structurally diverse set of aminodiols which give different relative orientations of the functional groups. The backbone segment amine function, also referred to as the reactive amino group, or amino combinatorial site, is protected with a base labile protecting group and one hydroxyl is blocked with a protecting group removable with mild acid such as a dimethoxytrityl (DMT) ether. The nitrogen protecting group is first removed from the backbone segment(s) and diverse functional groups added to the reactive amino group. Methods for achieving this are described below. A second monomer subunit can then be added through H-phosphonate coupling after removal of the acid labile protecting group on the terminal end of the compound on the solid support by, for example, treating with dilute acid. The intermediate hydrogen phosphonate diester can be oxidized with, for example, a solution of CCl_4 , Pyridine (1:1) containing 10 % (v/v) of a primary or secondary amine, resulting in the formation of a phosphoramidate linkage. Alternatively the H-phosphonate linkage can be oxidized to the phosphodiester or the phosphorothioate using standard methods and techniques. The second backbone segment can then be substituted with diverse functional groups as above. Any combination of backbone segments and functional groups at the backbone segment amino combinatorial sites can be introduced at any position of the oligomer and a large number of amines can be used at the oxidation step. By repeating these synthetic steps an oligomeric compound is synthesized having any desired sequence of functional groups on the monomer subunits, and amide derivatives at any

- 17 -

linking position. Random positions are introduced into the library by dividing the solid support into portions before the addition of the appropriate reagent. Either the monomers, the monomer substituents or the amines can be randomized. At least one position in the oligomer subsets must be fixed to allow iterative deconvolution. The library subsets are screened in the biological assay of interest. The most active subset defines the most active residue at the fixed position. Further rounds of synthesis and screening are used to determine the sequence of the most active compound in the library.

One feature of the present invention is the use of a nitrogen blocking group to block the reactive amino site. Once the first aminodiol monomer subunit is attached to the solid support, the nitrogen blocking group can be removed under basic (non hydrolytic) conditions. The nitrogen is then derivatized with the diverse functional group of choice. This group can be attached to the amino combinatorial site via a variety of linkage groups: amide, sulfonamide, carbamate, urea, aminoalkane, thiocarbamate, thiourea, etc. This can be accomplished by choosing the appropriate electrophile to derivatize the nitrogen. For example, carboxylic acids can be activated using peptide coupling reagents such as EDC, BOP or HATU. Other reagents which can be used include acid chlorides, -fluorides, -imidazolides, -anhydrides, sulfonyl chlorides, chloroformates, isocyanates, aldehydes (under reductive alkylation conditions), alkyl halides, isothiocyanates, etc. Thus each time a functional group is desired in a library it is introduced via the appropriate coupling conditions using simple starting materials.

The aminodiol monomer subunits in the combinatorial library each bear functional groups e.g. "letters" in addition to those that form linkages. When the aminodiol monomer subunits are linked together, these functional

groups provide diverse properties ("diversity") to the resulting oligomeric compounds. The functional groups include hydrogen-bond donors and acceptors, ionic moieties, polar moieties, hydrophobic moieties, aromatic centers, and electron-donors and acceptors. Together, the properties of the individual monomers contribute to the uniqueness of the oligomeric compounds in which they are found. Thus, a library of such oligomers would have a myriad of properties, i.e., "diversity." Collectively, the properties of the individual monomers that together form an oligomeric compound contribute to the uniqueness of such oligomeric compound and impart certain characteristics thereto for interaction with cellular, enzymatic or nucleic acid target sites.

A protecting group such as a member of the trityl family preferably can be used as the acid labile protecting group of one of the two hydroxyls of the aminodiol monomer subunit. The trityl family includes at least trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl. The dimethoxytrityl group is preferred and can be added by reacting the primary hydroxyl group with 4,4'-dimethoxytrityl chloride. Other hydroxyl protecting groups can be used. Representative hydroxyl protecting groups are described by Beaucage, et al., *Tetrahedron* 1992, 48, 2223. Preferred hydroxyl protecting groups are acid-labile, such as the trityl, monomethoxytrityl, dimethoxytrityl, trimethoxytrityl, 9-phenylxanthine-9-yl (Pixyl) and 9-(p-methoxyphenyl)xanthine-9-yl (MOX).

In other aspects of the present invention the use of acid labile groups which are stable to the trichloroacetic acid treatment used for DMT removal such as BOC-type protecting groups are used. They are stable to extended TCA treatment, but are removed by trifluoroacetic acid solutions (e.g. 5% in CH_2Cl_2). Another protecting group class which is compatible to this methodology is the allyl class. These

groups are cleaved using transition metal catalysts. These types of protecting group are particularly valuable in cases where the selective deprotection of a particular functional group is desired while the oligomer is still attached to the solid support, allowing a new reactive site to be uncovered. Additional protecting group tactics are possible: e.g. photolabile protecting groups are also compatible with this methodology.

In another aspect of the invention, nitrogen protecting groups that are stable to acid treatment and are selectively removed with base treatment are used to make reactive amino groups selectively available for substitution. Examples of such groups are the Fmoc (E. Atherton, R.C. Sheppard in *The Peptides*. S. Udenfriend, J. Meienhofer, Eds. Academic Press, Orlando, 1987, vol 9, p1-38), and various substituted sulfonyl ethyl carbamates exemplified by the Nsc group (V.V. Samukov, A.N. Sabirov, P.I. Pozdnyakov, *Tetrahedron Lett*, 1994, **35**, p7821; C.G.J. Verhart, G.I. Tesser, *Rec. Trav. Chim. Pays-Bas*, 1987, **107**, p. 621).

Heterocycles, including nitrogen heterocycles, suitable for use as functional groups include, but are not limited to, imidazole, pyrrole, pyrazole, indole, 1H-indazole, a-carboline, carbazole, phenothiazine, phenoxazine, tetrazole, thiazole, oxazole, oxadiazole, benzoxazole, benzimidazole, triazole, pyrrolidine, piperidine, pyridine, quinoline, piperazine and morpholine groups.

Purines and pyrimidines suitable for use as functional groups include adenine, guanine, cytosine, uridine, and thymine, as well as other synthetic and natural nucleobases (nucleosidic bases) such as xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 5-halo uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudo uracil), 4-

thiouracil, 8-halo, amino, thiol, thioalkyl, hydroxyl and other 8-substituted adenines and guanines, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine. Further purines and pyrimidines include those disclosed in United States Patent No. 3,687,808, those disclosed in the *Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, and those disclosed by Englisch et al., *Angewandte Chemie, International Edition* 1991, 30, 613.

Alkyl, alkenyl, and alkynyl groups according to the invention include but are not limited to substituted and unsubstituted straight chain, branch chain, and alicyclic hydrocarbons. Further, in the context of this invention, a straight chain compound means an open chain compound, such as an aliphatic compound, including alkyl, alkenyl, or alkynyl. A branched compound, as used herein, comprises a straight chain compound, such as an alkyl, alkenyl, alkynyl compound, which has further straight or branched chains attached to the carbon atoms of the straight chain. A cyclic compound, as used herein, refers to closed chain compounds, i.e. a ring of carbon atoms, such as an alicyclic or aromatic compound. The straight, branched, or cyclic compounds may be internally interrupted, as in alkoxy or heterocyclic compounds. In the context of this invention, internally interrupted means that the carbon chains may be interrupted with heteroatoms such as O, N, or S. However, if desired, the carbon chain may have no heteroatoms.

Further in the context of this specification aryl groups include but are not limited to substituted and unsubstituted aromatic hydrocarbyl groups. Aralkyl groups include but are not limited to groups having both aryl and alkyl functionalities, such as benzyl and xylyl groups. Preferred aryl and aralkyl groups include, but are not, limited to, phenyl, benzyl, xylyl, naphthyl, tolyl, pyrenyl, anthracyl, azulyl, phenethyl, cinnamyl, benzhydryl, and mesityl. These

- 21 -

can be substituted or unsubstituted.

The aliphatic and aromatic groups as noted above may be substituted or unsubstituted. In the context of this invention, substituted or unsubstituted, means that the compounds may have any one of a variety of substituents, in replacement, for example, of one or more hydrogen atoms in the compound, or may have no substituents. Typical substituents for substitution include, but are not limited to, hydroxyl, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, or alkyl, aryl, alkenyl, or alkynyl groups.

Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethers including polyethylene glycols, and other moieties known in the art for enhancing the pharmacodynamic properties or the pharmacokinetic properties. Typical conjugate groups include PEG groups, cholesterol, phospholipids, biotin, phenanthroline, phenazine, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes.

Metal coordination groups according to the invention include but are not limited to hydroxamic acids, catecholamide, acetylacetone, 2,2'-bipyridine, 1,10-phenanthroline, diacetic acid, pyridine-2-carboxamide, isoalkyldiamine, thiocarbamate, oxalate, glycol, histidyl and terpyridyl. Other metal coordination groups are known, as for example see Mellor, D.P., *Chemistry of Chelation and Chelating Agents in International Encyclopedia of Pharmacology and Therapeutics*, Section 70, The Chelation of Heavy Metals, Levine, W.G. Ed., Pergamon Press, Elmford, N.Y., 1979.

Solid supports according to the invention include controlled pore glass (CPG), oxalyl-controlled pore glass (see, e.g.,) or Poros --a copolymer of polystyrene/divinylbenzene.

Non-reactive functionalities used as functional groups,

- 22 -

such as groups that enhance pharmacodynamic properties, include groups that improve uptake and enhance resistance to enzymatic or chemical degradation. Non-reactive functionalities may also enhance pharmacokinetic properties.

5 In the context of this invention, such groups improve uptake, distribution, metabolism or excretion. Non-reactive functionalities include, but are not limited to, alkyl chains, polyamines, ethylene glycols, steroids, polyamides, aminoalkyl chains, amphipathic moieties, and conjugate
10 groups attached to any of the nitrogenous sites for attachment, as described above.


A number of functional groups can be introduced into compounds of the invention in a blocked form and subsequently deblocked to form a final, desired compound. In
15 general, a blocking group renders a chemical functionality of a molecule inert to specific reaction conditions and can later be removed from such functionality in a molecule without substantially damaging the remainder of the molecule (Green and Wuts, *Protective Groups in Organic Synthesis*, 2d
20 edition, John Wiley & Sons, New York, 1991). For example, amino groups can be blocked as phthalimido groups, as 9-fluorenylmethoxycarbonyl (Fmoc) groups, and with triphenylmethylsulfenyl, *t*-BOC or CBZ groups. Hydroxyl groups can be protected as acetyl groups. Representative
25 hydroxyl protecting groups are described by Beaucage *et al.*, *Tetrahedron* 1992, 48, 2223. Preferred hydroxyl protecting groups are acid-labile, such as the trityl, monomethoxytrityl, dimethoxytrityl, trimethoxytrityl, 9-phenylxanthine-9-yl (Pixyl) and 9-(*p*-methoxyphenyl)xanthine-
30 9-yl (MOX). Chemical functional groups can also be "blocked" by including them in a precursor form. Thus, an azido group can be used considered as a "blocked" form of an amine since the azido group is easily converted to the amine.

35 Additional functional groups according to the invention


- 23 -

include but are not limited to H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, carbocyclic alkyl, alkenyl or alkynyl or substituted carbocyclic, or aryl or substituted aryl where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, or alkynyl groups; an gluether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms; a nitrogen, sulfur or oxygen containing heterocycle; a metal coordination group; a conjugate group; halogen; hydroxyl (OH); thiol (SH); keto (C=O); carboxyl (COOH); amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R¹OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups.

Functional groups of the invention can be represented by structure:



where T is a single bond, a methylene group or a group having formula:



where:

- 24 -

R_{10} is =O, =S, or =NR₁₁;

R_5 and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;

each R_6 , R_7 , R_8 , R_9 , R_{11} , R_{12} and R_{13} are,

5 independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

10 p is 0 or 1;

q is 1 to about 10;

L is H, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted
 15 C₄-C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇ carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur
 20 containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)),
 25 nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂), sulfide (S-), disulfide (S-S), silyl, a
 30 nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and
 35 alkynyl groups.

03973334-03333

- 25 -

As used in this specification, a "functional" group is one that, when attached to a parent molecule, imparts to that molecule a particular and unique characteristic. It contributes diversity to the parent molecule by rendering the parent molecule different in some way from what it was before attachment of the group. Several chemical functional groups can be attached to a particular molecule and when considered together, the sum total of their properties will impart global diversity characteristics to the parent molecule. Each set of combinations of chemical functional groups on a particular molecule will modify the parent such that the parent molecule having each particular combinations of groups will be different from the parent molecule having any of the other combinations of groups. When all of the combinations of the groups on the parent are considered, a library of compounds will be formed that include all of the possible combinations of groups.

Oligomeric compounds of the invention can be synthesized with both the position and the choice of the chemical functional groups predetermined, or allowed to be selected by combinatorial selection. In the context of this invention, "combinatorial" does not mean arbitrary, haphazard or indiscriminate. In the context of this invention, "combinatorial" is construed to mean that within the totality of the population of oligomeric compounds that can be formed using a particular set of functional groups and a particular location of combinatorial sites within the oligomeric compound, there will be sub-populations of each of the possible species. Thus, each of the different combinations of a) choice of functional group and b) positioning of the functional groups will be represented.

"Combinatorial" is distinct from "random." To illustrate the distinction, if all or nearly all possible combinations are present in the total molecular population, then it is a combinatorial population of molecules. If,

- 26 -

however, only one or a small number of molecules from that total population is selected, then the selected molecule or molecules might be randomly selected if it is picked at whim or will from the total population. When the totality of the population is considered, all species are present and it is not a random population. If a systematic selection was made until the totality of the population was exhausted, then all of the species would eventually be selected, however, the order of selection might be random. Thus, in certain preferred embodiments, a pre-ordered selection and/or location of chemical functional groups will be present. In further preferred embodiments, a combinatorialized population of all possible combinations and ordering of the chemical functional groups is present. In even further preferred embodiments, the sequence is modulated between fixed and combinatorial. This is especially useful, as for example, in certain deconvolution strategies.

"Deconvolution" is construed to mean taking the totality of a population and systematically working through that population to establish the identity of a particular member, selected members, or all members of the population. In deconvoluting a combinatorial library of compounds, systematic selection is practiced until an individual oligomeric compound or a group of individual oligomeric compounds having a particular characteristic, as for instance being an active species in a specific functional assay, is identified.

Conjugate groups can also be used as functional groups in the present invention. Conjugate groups include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterols, phospholipids, biotin, phenanthroline, phenazine,

phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA or protein targets. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve compound uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992, United States Patent Application Serial No. 116,801, filed September 3, 1993, and United States Patent No. 5,218,105. Each of the foregoing is commonly assigned with this application. The entire disclosure of each is incorporated herein by reference.

Amines include amines of all of the above alkyl, alkenyl and aryl groups including primary and secondary amines and "masked amines" such as phthalimide. Amines of this invention are also meant to include polyalkylamino compounds and aminoalkylamines such as aminopropylamines and further heterocycloalkylamines, such as imidazol-1, 2, or 4-yl-propylamine.

Amino groups amenable to the present invention have the formula $N(Y_0)T_0$, wherein:

Y_0 is H, or $[Q_2]_j-Z_2$;

T_0 is $[Q_1]_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently, are C_2-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_4-C_7 carbocyclo alkyl or alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7-C_{14} aralkyl;

j and k independently, are 0 or 1;

Z_1 and Z_2 independently are H, C_1-C_2 alkyl, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, C_6-C_{14} aryl, C_7-C_{15} aralkyl, halogen,

CH=O, OR₁₂, SR₁₂, NR₁₂R₁₃, C(=NH)NR₁₂R₁₃, CH(NR₁₂R₁₃),
NHC(=NH)NR₁₂R₁₃, CH(NH₂)C(=O)OH, C(=O)NR₁₂R₁₃, C(=O)OR₁₂, a metal
coordination group, a reporter group, a nitrogen-containing
heterocycle, a purine, a pyrimidine, a phosphate group, a
5 polyether group, or a polyethylene glycol group.

In some preferred embodiments, the oligomeric compounds
of the invention are from 2 to about 30 aminodiol monomers
in length. More preferred embodiments have from 2 to about
15 aminodiol monomers, and especially preferred embodiments
10 have from 2 to about 6 aminodiol monomers.

To synthesize a combinatorial library having a large
degree of chemical diversity is an important aspect of the
present invention. Chemical diversity is introduced at one
level by varying the nature of the phosphorus linkage.
15 Phosphorus linkages amenable to the present invention
include phosphodiester (OPO), phosphorothioate (OPS), and
phosphoramidate (OPN). The desired combinatorial library
can be prepared with a single type of phosphorus linkage, or
with different linkages at each position of the oligomer.
20 For example, a single OPS linkage can be selectively
introduced at any position in a OPO oligomer. In fact, all
possible combinations of OPO, OPS, and OPN linkages can be
introduced selectively into the oligomeric compounds. The
presence or absence of a type of linkage at a particular
25 position in an oligomer will have profound effects on the
properties of the molecule.

In the case of phosphoramidate linked libraries, a
further level of diversity is possible by oxidizing the H-
phosphonate diester linkage with a solution such as CCl₄,
30 Pyridine (1:1) containing 10 % (v/v) of a primary or
secondary amine, resulting in the formation of a
phosphoramidate linkage. Any of the H-phosphonate monomer
subunits can be introduced at any position of the oligomer
and a large number of amines can be used at the oxidation
35 step. Thus it is possible to introduce a wide variety of

- 29 -

amines into the oligomeric compound at H-phosphonate linkages by oxidation to the OPN linkage. It is possible to have the same amine substituents at each OPN linkage or a different amine at each position. In the preparation of a combinatorial library from a set of monomer subunits, amines and function groups, all possible combinations are synthesized simultaneously.

Chemical diversity can be generated at several levels in SURFTM libraries. We have described below the preparation of a number of monomer subunits. These monomers have been prepared to explore two aspects of chemical diversity: first a wide number of functional groups are available, covering a range of chemical properties. Second, these functional groups can be attached directly to combinatorial sites or via tethering groups to combinatorial sites. Many different tethering groups are commercially available and are amenable to the present invention. The use of tethering groups alters the presentation of functional groups, in space in different ways, allowing variable flexibility.

Hydrogen phosphonate chemistry allows additional chemical modifications to be introduced into oligomeric compounds. Oligonucleotide phosphodiester and phosphorothioates have been prepared using this approach, (see Froehler, B. C., Matteucci, M. D. *Tetrahedron Lett.* 1986, 27, 469-472), as well as oligonucleotide phosphoramidates (see Froehler, B. C. *Tetrahedron Lett.* 1986, 27, 5575-5579. Letsinger, R. L., Singman, C. N., Histan, G., Salunkhe, M. *J. Am. Chem. Soc.* 1988, 110, 4470-4471. The synthesis of oligomeric compounds containing both phosphodiester and phosphoramidates was reported, as well as the use of phosphoramidite chemistry in conjunction with the synthesis of phosphoramidates (see Jung, P. M., Histan, G., Letsinger, R. L. *Nucleosides & Nucleotides*, 1994, 13, 1597-1605). In this latter work, alternating phosphodiester and phosphoramidate oligomeric compounds were prepared by

- 30 -

coupling phosphoramidites and H-Phosphonates to the growing oligomer, followed by the appropriate oxidation step. In general, however, all the examples described thus far have incorporated the same amine substitution at all

5 phosphoramidate linkages in the oligomer. These studies have shown the feasibility of using the phosphoramidate bond as an additional site for the incorporation of diverse functional groups.

10 A wide variety of amines can be used in the oxidative step, and the monomers of the present invention support the necessary chemistry. Thus, for the preparation of combinatorial libraries incorporating phosphoramidate linkages, the monomer subunits of the present invention used as the corresponding H-Phosphonate monoesters. In one aspect of the
15 present invention this is accomplished using PCl_3 and imidazole as the phosphitylating reagent (see Garegg, P. J., Regberg, T., Stawinski, J., Strömberg, R. *Chem. Scr.* 1986, 26, 59-62). These H-phosphonates monomer subunits may be oligomerized on solid support by activation with pivaloyl
20 chloride, adamantoyl chloride or other appropriate activating agent. The intermediate H-Phosphonate diesters are oxidized to the phosphate diesters in high yields using iodine in aqueous pyridine. This allowed for the comparison of the coupling efficiency of the H-phosphonate and
25 phosphoramidite methods. Essentially the same coupling efficiency is achieved with both methodologies. The H-phosphonate diesters are converted to phosphoramidates by the use of a 10% solution of the appropriate amine in pyridine/ CCl_4 (1:1). Under these conditions, a H-phosphonate
30 diester is oxidized to a phosphoryl chloride via an Arbuzov reaction, followed by displacement of the chloride by a primary or secondary amine. The second step has proven to be quite general, with a wide variety of amines giving satisfactory yields. Moreover, the yield of phosphoramidate
35 is comparable to the yield of phosphodiester.

- 31 -

Several types of libraries are available through this methodology. The simplest kind is a library made from a set of monomer subunits of the present invention, and a set of 4 to 16 or more functional groups, of 2 to 10 or more monomer subunits in length, which is substituted at phosphorus with a single amine. These libraries are prepared by split bead synthesis, following the H-phosphonate synthesis protocol rather than phosphoramidite chemistry.

In one aspect of the present invention, intermediate H-phosphonate diesters are left intact until the final step. At that point the oligomer library pools are oxidized with CCl_4 /Pyridine containing 10% of the appropriate primary or secondary amine. The library therefore is composed of all possible sequences of the monomers, separated into subsets unique at a fixed position, linked together by a constant phosphoramidate linkage. It should be evident that the final properties of the library will be determined by the choice of amine used in the oxidation step. Thus, water solubility, pharmacokinetics and pharmacodynamics of the library components can be modulated by the choice of amine.

It is also possible to prepare oligomer libraries with mixed linkages by having an intermediate oxidation step (see Gryaznov, S. M., Sokolova, N. I. *Tetrahedron Lett.* **1990**, 31, 3205-3208; Gryaznov, S. M., Potapov, V. K. *Tetrahedron Lett.* **1991**, 32, 3715-3718; Farooqui, F., Sarin, P. S., Sun, D., Letsinger, R. L. *Bioconjugate Chem.*, **1991**, 2, 422-426; Iso, Y., Yoneda, F., Ikeda, H., Tanaka, K., Fuji, K. *Tetrahedron Lett.* **1992**, 33, 503-506). Thus, a portion of the oligomer library is synthesized by H-phosphonate chemistry, which can be oxidized with (R_2NH , CCl_4/Py or S_8 , CS_2/TEA or H_2O , CCl_4/Py), and a second portion of the library synthesized and oxidized with a second set of reagents. This creates a chimeric library, where a segment of the random oligomeric compounds in each subset has a different linkage

- 32 -

than the rest of the molecule.

By extension of this methodology, it is possible to incorporate a different linkage at each position of the oligomer library by having a different oxidation step after each monomer subunit coupling. The linkage can be combinatorialized by performing a separate oxidation on a portion of the H-phosphonate diester-linked solid support, followed by pooling of the subsets in the same way that the monomer subunit positions are randomized. Thus, each monomer and the linkage between them can be randomized by a split synthesis strategy.

One advantage of the present invention is that the simple design of monomer subunits of the inventions allows for combining rational drug design with screen mechanisms for thousands of compounds. This is achieved by using the monomer subunits of the invention in a combinatorial techniques such as the SURF™ strategies.

In one preferred embodiment, functional groups appended to oligomeric compounds of the invention are selected for their potential to interact with, and preferably inhibit, the enzyme PLA₂. Thus, the oligomeric compounds of the invention can be used for topical and/or systematic treatment of inflammatory diseases including atopic dermatitis and inflammatory bowel disease. In selecting the functional groups, advantage can be taken of PLA₂'s preference for anionic vesicles over zwitterionic vesicles. In selecting the backbone segments that bear these functional groups, further advantage can be taken of fact that the natural substrate of PLA₂ contains a phosphate group. Therefore, phosphodiester or phosphorothioate and other phosphate linked oligomeric compounds may be selected, providing a negatively charged compound for binding with the positively charged interfacial binding site of PLA₂.

Compounds of the invention also include aromatic functional groups to facilitate binding to the cleft of the

- 33 -

PLA₂ enzyme. (see, Oinuma, et al., *J. Med. Chem.* **1991**, *34*, 2260; Marki, et al., *Agents Actions* **1993**, *38*, 202; and Tanaka, et al., *J. Antibiotics* **1992**, *45*, 1071). Benzyl and 4-hexylbenzyl groups are preferred aromatic groups. The compounds of the invention can further include hydrophobic functional groups such as tetraethylene glycol groups. Since the PLA₂ enzyme has a hydrophobic channel, hydrophobicity is believed to be an important property of inhibitors of the enzyme.

10 In certain embodiments of the invention, aminodiol monomer subunits are incorporated into libraries of oligomeric compounds and increasingly less complex subsets of oligomeric compounds are identified in combinatorial screening techniques such as the above described SURF™ technique by successive rounds of screens. The PLA₂ assay can be effected using a combinatorial screening strategy such as the SURF™ strategy. For this assay, the libraries of oligomeric compounds are screened for inhibition of human type II PLA₂ enzymatic activity. Typically, these libraries contain about 100 to 100,000 different compounds. Successive iterations of the SURF™ technique is effected to select unique oligomeric compounds from the library. The libraries additionally can be screened in other *in vitro* assays to determine further mechanisms of inhibition.

25 Upon identification of oligomeric compounds in a first phase of screening, further modifications can be made to the contents of the libraries of oligomeric compounds. For example, if a first iteration of screening results in an active compound that contains a benzyl group, then in subsequent iterations of the screen this aromatic residue can then be varied using substituted benzyl groups. In this way, structural activity is identified in a stepwise manner to define potent inhibitors of the enzymatic activity.


To maximize the identification of a tight binding oligo-

- 34 -

meric inhibitor of PLA₂ via a combinatorial approach, an array of functional groups typically are included in a randomized library.

Aminodiol monomer subunits can be linked with one another to form homopolymeric structures or they can be linked with nucleotides and/or other moieties to form chimeric structures. For example, chimeric structures can be formed that include one or more regions or "stretches" of the monomer subunits of the invention joined to one or more regions or "stretches" of naturally occurring or synthetic oligonucleotides or to other synthetic or natural oligomeric compounds such as peptides, peptoids, peptide nucleic acids, oligo and/or polysaccharides. Further, oligomeric compounds of the invention can be incorporated into chimeric structures along with the compounds disclosed in the patent application entitled "Monomeric Diols And Phosphate Linked Oligomeric compounds Formed Therefrom," Serial No. 08/179,970 filed January 11, 1994; which is the parent case for PCT/US95/0049, filed January 11, 1995; and the patent application entitled "Oligonucleotide Mimics Having Nitrogen-Containing Linkages," Serial No. 08/180,124, filed January 11, 1994; which is the parent case for PCT Application bearing attorney docket ISIS-1852, filed January 11, 1995. The foregoing patent applications are commonly assigned, and are incorporated herein by reference.

In one aspect of the invention, a combinatorial library of oligomeric compounds is synthesized by first attaching a plurality of aminodiol monomer subunits, each having structure I, II, III, IV, V, VI, VII, VIII, IX, X, or XI, below, to a solid support.



The first aminodiols monomer subunits have base labile protecting groups for R_1 and a hydrogen or an alkyl group for R_2 , if present. One of R_3 and R_4 is an acid labile protecting group and the other is hydrogen e.g. a free hydroxyl. The free hydroxyl is reacted with a group on the solid support to effect the attachment of the aminodiols monomer subunit to the solid support. Alternatively other bifunctional groups such as ethylene glycol are attached to the solid support with many such compounds known in the art and many commercially available. Solid supports are also available with linking groups previously attached, ready for use

- 36 -

without a derivatization step.

The base labile protecting group on the reactive amino group is removed by contacting with a base such as 10% piperidine in DMF or DBU (diazabicycloundecene) in pyridine, and a substituted carboxylic acid or another electrophilic reagent (acid chloride, sulfonyl chloride, etc.) is then covalently linked to the backbone segment amino combinatorial site using standard coupling methods (for examples see Bodansky, M., *Principles of Peptide Synthesis*, 1984, Springer-Verlag, Berlin). Carbamates can be obtained by the treatment of the amine with an appropriate alkyl or aryl chloroformate. Carbamates can be obtained by the treatment of the amine with an appropriate alkyl or aryl chloroformate, in the presence of a catalyst such as pyridine. A urea or thiourea can be formed by reacting the backbone segment amino combinatorial site with an isocyanate or isothiocyanate, or by treatment with carbonyl diimidazole followed by a second amine, in the presence of base. Sulfonamides can be prepared from the amine by the reaction with a sulfonyl chloride in the presence of a base. The nitrogen can be alkylated by treatment with a halide such as the illustrative halides (benzyl bromide, 3-methylbenzyl bromide, 3-methoxybenzyl bromide or 3-nitrobenzyl bromide) used in the examples below. A wide spectrum of halides can be used for this purpose. Additionally, amino compounds can be functionalized by reaction with an aldehyde or ketone forming a Schiff base. The Schiff base is then reduced in the presence of a reducing agent such as NaCNBH_3 .

Functional groups that require protection are derivatized using acid labile protecting groups which are stable to TCA. The acid labile hydroxyl protecting group is removed by treating with an acid (3% TCA in dichloromethane) and the resulting free hydroxyl is reacted with a second aminodiol monomer subunit H-phosphonate having a base labile protecting group for R_1 and a hydrogen or an alkyl group for

- 37 -

R₂, if present. One of R₃ and R₄ is an acid labile protecting group and the other is an activated phosphite. The free hydroxyl of the first monomer subunit on the solid support is reacted with the activated phosphite of the second monomer subunit to form a phosphite linkage thereby increasing the length of the growing oligomeric compound by one monomer subunit.

The phosphite linkage is oxidized to a phosphoramidate using a 10% solution of a primary or secondary amine in CCl₄/pyridine. The base labile protecting group on the second monomer subunit is then removed with piperidine as above. The resulting amino group is further reacted with an activated functional group. If desired, additional monomer subunits can be added by repeating the synthetic steps described above.

Monomer subunits of the invention can be used to prepare oligomeric compounds having either preselected sequences or sequences determined via combinatorial strategies. One useful combinatorial strategy is the above-noted SURFTM strategy, which is disclosed and claimed in United States Patent Application Serial No. 749,000, filed Aug. 23, 1991, and PCT Application US92/07121, filed Aug. 21, 1992, both of which are commonly assigned with this application. The entire disclosure of these applications are herein incorporated by reference.

The following is an example of the methods used for the synthesis of a simple combinatorial library. The variable parameters which can be controlled are the following: the backbone segments, the functional groups attached to the backbone segment amino combinatorial site, the amines attached at the phosphoramidate linkage, and the length of the oligomer. In the example described here a very simple library composed of only 32 molecules will be described. This library is prepared by using two different backbone segments, A and B, two carboxylic acids X and Y, and two

- 38 -

amines, P and Q. 1) One begins by weighing equal amounts of solid supports derivatized with A and B. The two solid supports are mixed together by suspending the gels in DMF/dichloromethane (1:1) or other solvent, and mixing gently. If more than two backbone segments are used in the library, all the different supports are mixed together at this stage. The mixture of supports is then treated with piperidine in DMF to remove the N-protecting group from both supports. 2) The mixture is separated into a number of equal portions corresponding to the number of reagents to be added in the next step, in this case two. To the first mixture is added activated carboxylic acid X, to the is added activated Y. Once the reaction is complete, the solid supports are mixed as above to give an equimolar mixture of all four possible combinations. The mixture is then treated with trichloroacetic acid to remove the DMT protecting groups on the monomer subunits and provide a free OH group. 3) The mixture is separated into a number of equal portions corresponding to the number of monomer subunits to be added, in this case two. To each mixture a monomer subunit H-phosphonate is then coupled, the solid supports are again mixed, and the support is again divided into equal portions corresponding to the number of amines to be added, in this case two. 4) At this stage all eight possible combinations are present. Each portion is treated separately with a solution an amine, P or Q in CCl₄/Pyridine. The portions are then pooled, treated with piperidine in DMF and split. 5) Each mixture of 16 compounds on solid support treated with an activated carboxylic acid, X and Y. The result is two unique pools of 16 compounds each, which can then be deprotected to remove the terminal DMT group and any protecting groups on the functionalities, and cleaved from the solid support. It is also possible to add further combinatorial sites by coupling additional monomer subunits and extending the library molecules. We have used the

- 39 -

following nomenclature to describe mixtures of compounds in a concise manner: X represents a fixed position, that is the backbone segment or functional group on the amino combinatorial site at that position is unique and serves to define the pool composition. N represents an equimolar mixture of all possible structures at a particular position. The subscript defines the synthetic step in the library synthesis. In the above example the library made was $X_5N_4N_3N_2N_1$, where $X_5 = X$ or Y , N_4 is an equal mixture of P and Q, $N_3 = A$ and B , $N_2 = X$ and Y , and $N_1 = A$ and B . In this case the fixed position is the last one, but similar methods can be used to fix any position in the library independently. It is also possible to have a single structure or functionality at any position. Once the most active moiety at position 5 is determined for a particular assay e.g. A, then two additional pools are synthesized having this residue in all cases. A second position is fixed and the remainder randomized: for example $A_5X_4N_3N_2N_1$, where N_{1-3} are as above and X_4 is either P or Q. A unique structure is determined after 5 rounds of synthesis and screening. The advantage is uncovered when more than two possible components are used in each position: If ten different components are available at each position then 100,000 unique structures exist, yet only 5 rounds of synthesis and screening are necessary.

Illustrative of the SURF™ strategy is a 2'-O-methyl oligonucleotide library (see, Ecker et. al., *ibid.*) shown in Table I, below. Table I describes the selection of a 2'-O-methyl oligonucleotide for binding to an RNA hairpin. The K_D 's, i.e., the binding constants, were determined by gel shift. "X" is used to indicate the position being varied and underlining is used to indicate positions that become fixed during successive iterations of the SURF™ strategy.

TABLE I

- 40 -

		K _p (mM)			
		X=A	X=C	X=G	X=T
5	Round 1				
	NNNNXNNNN	22	<u>10</u>	>100	>100
	Round 2				
	NNNNCNXNN	>10	<u>4</u>	>10	>10
	Round 3				
	NNXNCNCNN	>10	<u>0.5</u>	>10	>10
10	Round 4				
	NNCXCNXNN	>10	<u>0.15</u>	>10	>10
	Round 5				
	NNCCCXCNN	<u>0.08</u>	>1	0.4	>1
15	Round 6				
	NNCCCACXN	<u>0.05</u>	>0.5	0.08	>0.5
	Round 7				
	NXCCCACAN	>0.1	>0.1	<u>0.03</u>	>0.1
	Round 8				
	NGCCCACAX	0.05	<u>0.02</u>	0.05	0.04
20	Round 9				
	XGCCCACAC	0.03	0.05	0.02	<u>0.01</u>

One aspect of the present invention is the inclusion of monomer subunits of the invention in the above-described SURF™ strategy. The SURF™ strategy is equally applicable to libraries of chemical compounds of the present invention in a completely parallel manner. Many other assays are also used as the selection criteria to deduce a winning sequence with the highest activity. The functional groups appended to the reactive amino groups of the oligomeric compounds of the invention can be of various structures that impart particular interactive properties to the oligomeric compounds. These chemical functional groups can effect interactions of at least the following types: hydrogen-bond donors and acceptors, ionic, polar, hydrophobic, aromatic,

- 41 -

electron donors and acceptors, pi bond stacking or metal binding. As a result of such interactions, the oligomeric compounds of the invention have unique properties effecting the overall global shape, the conformational space, electron density, dipole moment and ability of the compound to interact with enzyme pockets and other binding sites and other similar properties.

To detect an active sequence generated via a combinatorial technique, the concentration of the active molecule is selected to be of sufficiently great that the molecule can be detected within the sensitivity of the chosen assay. As will be recognized, the number of unique structures within a subset produced via a combinatorial technique depends on the length of the oligomer and the number of different functionalities employed. The number of structures is given by the product of the number components at each variable position. This is illustrated in Table II. Table II also indicates the concentration of each sequence when the subset concentration is 100 μ M, a typical high-test concentration. We have found that the complexity of the library can be based upon an estimate of the expected IC_{50} (i.e., a concentration at which 50% of enzyme activity is inhibited) that is desirable in a final oligomeric compound. For an expected IC_{50} of 100 nM, the complexities shown in Table II are acceptable, that is, the libraries shown in Table II have complexities that would allow detection of a unique sequence with an IC_{50} of about 100 nM or less.

TABLE II Complexity of Libraries

	Variable Positions (Mer)	Sequences nM Each Sequence	
		Per Subset	At 100 μ M Subset
	5 Components		
35	4-mer	125	800

- 42 -

	5-mer	625	160
	6 Components		
	4-mer	216	463
	5-mer	1,296	77
5	7 Components		
	4-mer	343	291
	8 Components		
	4-mer	512	195
	10 Components		
10	4-mer	1,000	100

The functional groups or components can also be referred to as "letters." The use of such terminology reflects the fact that the different functional groups on the compounds of the invention are positioned in sequences (either predetermined or by random selection) much like letters of the alphabet, hence the term "letter." These letters can be "reactive" or "non-reactive." By "reactive," it is meant that they will interact with a target molecule in some manner (that need not but can be predefined). By non-reactive," it is meant that they are not designed to primarily interact with a target molecule, and in fact while they may interact with the target molecule, the primary purpose of the non-reactive moieties are to impart other properties to the molecule such as, but not necessarily limited to, effecting up-take, distribution, metabolism or identification.

A further advantage of this invention is the ability to synthesize oligomeric compounds that, in addition to or in place of variability in the sequences of the diverse functional groups, have an asymmetric sequence of monomer subunits. Stated otherwise, the monomer subunits can also vary within the oligomeric compounds. This is easily accomplished by using different aminodiol monomer subunits


















- 43 -

that eventually become the backbone of the oligomeric compounds.


The oligomeric compounds of the invention can be used in diagnostics, therapeutics and as research reagents and kits. They can be used in pharmaceutical compositions by including a suitable pharmaceutically acceptable diluent or carrier. In preferred embodiments, the compounds of the invention act as inhibitors of enzymes such as phospholipase A₂; as inhibitors of pathogens such as virus, mycobacterium, bacteria (gram negative and gram positive), protozoa and parasites; as inhibitors of ligand-receptor interactions such as PDGF (platelet derived growth factor), LTB₄ (leukotriene B₄), IL-6 and complement C5_A; as inhibitors of protein/protein interactions including transcription factors such as p50 (NF- κ B protein) and fos/jun; for the inhibition of cell-based interactions including ICAM induction (using inducers such as IL1- β , TNF and LPS) and as herbicides and insecticides. In other preferred embodiments, the compounds of the invention are used as diagnostic reagents for each of the above noted biological entities, and as reagents in assays and as probes. In other preferred embodiments, the compounds of the invention are used to chelate heavy metals and as imaging agents.

The functional groups can be selected based on chain length, i.e. short versus long, based on the use of a ring versus a linear group, use of an aromatic versus aliphatic group, use of a functionalized group versus a non-functionalized group, to mention only a few of the wide variety of chemical functional groups available. Indeed simply varying functional moieties, e.g. acid, alcohol, aldehyde, amide, amine, amidine, azo, azoxy, double bond, ether, ethylene oxide, guanidine, halide, haloalkyl, hydrazine, hydroxylamine, ketone, mercaptan, nitrate, nitrile, nitro, nitroso, quaternary nitrogen, sulfide, sulfone, sulfoxide, triple bond, urea, etc. on a single

backbone segment amino combinatorial site, e.g. a simple alkyl group, yields a vast array of diversity functions. When this is expanded to include placement of such varied functional moieties on a broad platform of backbones, e.g. a series of alkyl compounds, a series of aryl compounds, a series of alicyclic compounds, etc., the potential for a vast array of chemical functional groups is apparent. Other chemical functional groups include alkyl, alkenyl, alkynyl, alicyclic and substituted alkyl, alkenyl, alkynyl, alicyclic; aryl and substituted aryl; aralkyl, substituted aralkyl, heterocycles, nucleobases such as pyrimidines and purines, metal chelating groups and moieties as found in the α -position of amino acids, such as those shown below:



- 45 -



The various combination of reactions that can be effected utilizing the teaching of this invention can be further demonstrate by use of a logical flow scheme. This flow scheme is set forth in Figures 1A and 1B.

5 Referring to the figures, at the start 10 of the synthesis, an aminodiol monomer subunit or group of aminodiol monomer subunits is/are selected. The selected monomer or monomers are processed at process step 12 wherein the aminodiol monomer subunit or subunits is/are attached to a
 10 solid support as per Examples 26, 30, 35, 45, 46, 57, 55, and 61, set forth below. The aminodiol monomer subunit may be attached to the solid support via a H-phosphonate diester linkage, through a succinyl linkage, or through an ethylene-glycol linkage. Many other types of linkages are known in
 15 the art and are amenable to the present invention. A group of aminodiol monomer subunits may be advantageously attached at this step by utilizing a split bead synthesis. The process steps effected at process step 12 can be otherwise characterized as a selecting and attaching procedure depicted
 20 by box 102.

From the procedure box 102, the path of the flow scheme intersects convergent point 14. Convergent point 14 represents the point in the iterative process of the invention wherein growing oligomeric compounds from later
 25 steps are reintroduced via this iterative process to undergo further oxidation, amine group functionalization and elongation as discussed below.

Downstream of the convergent point 14, the processes of the invention further include an oxidization step effected
 30 via procedures included in the general oxidization procedure depicted by box 104. At decision block 16, within the

- 46 -

oxidization procedure generally depicted at box 104, the product of process step 12 (or the growing oligomeric compounds introduced at convergent point 14) is (are) either converted by oxidation from P^{III} to P^V compounds via the positive branch of this step or the P^{III} phosphite linkages are not oxidized and the compounds are maintained with P^{III} linkages and are taken to the next decision step via the negative branch of decision block 16 leading to convergent point 42. For those products of process step 12 (or the growing oligomeric compounds introduced at convergent point 14) wherein oxidization is desired during this iteration of the process (those moving along the positive branch of decision block 16), at decision block 18 they are either converted to phosphoramidates via the positive branch leading from decision block 18 or they are converted into phosphorothioates or phosphodiester via the negative branch leading from decision block 18.

At decision block 20, growing oligomeric compounds are either oxidized to phosphodiester at process step 24 via the positive branch of decision block 20 or they are oxidized to phosphorothioates at process step 25 via the negative branch of decision block 20. Oxidation to either the phosphorothioate or phosphodiester, as illustrated in Example 69, can be effected simultaneously on multiple positions on the same growing oligomeric compound, as for instance if the negative branch was taken at decision block 16 for one or more previous iterations of the process. If there are previous unoxidized sites and if the positive branch of decision block 16 is selected during the current iteration, multiple monomeric units are oxidized simultaneously. The resulting phosphodiester and phosphorothioates converge at convergent point 38 and from there lead to convergent point 40.

Following the positive branch from decision block 18, growing oligomeric compounds at decision block 22 can be

- 47 -

converted to phosphoramidates by one of three process steps. First, the decision to effect substitution using a single amine is effected at decision block 22. Growing oligomeric compounds following the positive branch from decision block 22 are converted at process step 26 using a single amine in solution as illustrated in Example 67, to give phosphoramidate linkages having uniform substituents.

Growing oligomeric compounds following the negative branch from decision block 22 are combinatorialized by one of two processes. Growing oligomeric compounds following the negative branch from the competitive combinatorialization decision block 28 are combinatorialized at process step 30 by a split bead synthesis. Growing oligomeric compounds following the positive branch of decision block 28 are combinatorialized at process step 32 using a mixture of amines as illustrated in Example 67 Method B.

Growing oligomeric compounds that may have been oxidized (via the positive branch of decision block 16) or may not have been oxidized (via the negative branch of decision block 16) converge at convergent point 42 within the general oxidization procedure box 106. From convergent point 42 the process is continued on Figure 1B via connector point 44 that connects Figure 1A and Figure 1B. A further connector point, point 46, leads back from Figure 1B to Figure 1A.

At decision block 48 within the generalized functionalization procedure depicted by box 106, following the positive branch of decision block 48, the growing oligomeric compounds, at process step 50, are deblocked to expose a free amino site on the growing oligomeric compound. This free amino site is now available for functionalization.

Alternatively, following the negative branch from decision block 48, amino deblocking and functionalization is bypassed. Following the negative branch of decision block 48, the amine protecting groups are maintained and the growing oligomeric compound is taken directly to convergent point 66.

- 48 -

If in one or more earlier iterations of the process, the negative branch of decision block 48 was selected, there will be multiple amino sites that now are protected and can be deblocked for functionalization. Thus functionalization can be effected at one or at more than one site during any one iteration of the functionalization procedure generally depicted by box 106.

Following the positive branch of the functionalization procedure decision block 48, after deblocking at process step 50, several alternatives are available for functionalization of the resulting free amino site(s). At decision block 52, functionalization with a single functional group or with multiple functional groups is queried. Following the positive branch from decision block 52, functionalization is effected with a single functional group. This functionalization can be performed on a single position or, as noted above, if the functionalization procedure was bypassed in prior iterations of the process, multiple amino positions in a growing oligomeric compound or combinatorial library of such compounds can be functionalized simultaneously. The negative branch of decision block 52 leads to decision block 56 where the deprotected amino site on the growing oligomeric compound can be functionalized with multiple reagents via two different combinatorialization procedures. Following the negative branch from decision block 56, combinatorialization of the free amino site(s) of the growing oligomeric compound(s) is effected at process step 58 using a split bead synthesis. Following the positive branch from decision block 56, combinatorialization of the free amino site(s) is effected at process step 60 by using a mixture of compounds e.g. carboxylic acids or acid halide as, for example, via Examples 63 and 65.

The products of process steps 58 and 60 converge at convergent point 62 and these in turn converge with the products of process step 54 at convergent point 64. Both of

- 49 -

these points converge at convergent point 66 where intermediates that were shunted by the deblocking process step 50 via the negative branch of the decision block 48 also converge.

5 A query is made at decision block 68 to discontinue synthesis or to further elongate the growing oligomeric compound. If the current oligomeric compounds is of sufficient length, at decision block 68, a decision to effect final deblocking, generally depicted by box 108, is made.
10 Following the negative branch of decision block 68, final deblocking is effected at process step 72 to give the final product or products 78.

 If further extension of the growing oligomeric compound is desired, the positive branch of decision block 68 is
15 followed. This leads to the extension procedure generally depicted by box 110. To effect oligomeric compound elongation, at process step 70 a hydroxyl protecting group is removed and at process step 74 a further aminodiol monomer subunit is added. The growing oligomeric compound is now re-
20 introduced at conversion point 14 (via connecting point 46) into the process loop for a further iteration of synthesis depicted in Figure 1A. Here the process of selection of paths independently leading to or bypassing oxidation and/or functionalization is then repeated.

25 The process of the invention can generally be characterized as first selecting and attaching one or more monomer subunits to a support. This corresponds generally to procedure box 102 of the figures. For the moment bypassing the oxidization procedure, generally depicted by box 104, and
30 the functionalization procedure, generally depicted by box 106, the hydroxyl blocking group of the monomeric subunits can be removed and a further iteration of the addition of monomer units effected. This is accomplished by the extension procedure generally depicted by box 110.

35 If during this iteration of the process, oxidization of

- 50 -

the phosphate linker is desired, oxidization will be effected via the oxidization procedure generally depicted by box 104. Further if prior to the next iteration but post the oxidization procedure, functionalization of the amino group of the last monomer subunit(s) added is desired, functionalization is effected via the functionalization procedure generally depicted by box 106.

If the growing oligomeric compound or compounds is/are of sufficient length, synthesis is halted and the compound or compounds are deblocked and cleaved from their supports via the final deblocking procedure, generally depicted by box 108.

It is of course recognized that the oxidization step of the first monomer subunit can be effected or oxidization of each subsequently added monomer subunit can be effected during the iteration of the process when they are added. Alternatively, oxidization need not be effected for each monomeric subunit added but can be effected during some subsequent iteration of the process. It is further recognized that functionalization of the amino site can be effected for the first monomer subunit and it can be effected for each added monomer subunit. Alternatively, functionalization need not be effected for each monomeric subunit added but can be effected during some subsequent iteration of the process.

EXAMPLE 1

N-Fmoc-*trans*-4-Hydroxy-L-Proline

trans-4-Hydroxy-L-proline (5.00 g, 38.2 mmol) and NaHCO₃ (8.00 g, 95.2 mmol) were suspended in 150 ml H₂O/Dioxane (1:1). Fluorenylmethyl chloroformate (11.4 g, 44.0 mmol) in 25 ml toluene was added dropwise. The temperature of the reaction was not allowed to rise above 25 °C during the addition. The mixture was stirred vigorously overnight, and

- 51 -

then quenched with 50 ml saturated NaHCO₃ solution and 50 ml water. The solution was then extracted with 100 ml diethyl ether. The aqueous layer was acidified to pH 1 with concentrated HCl, and extracted twice with ethyl acetate, and the organic extracts washed with brine. The solution was dried with MgSO₄, filtered and the solvent removed in vacuo. The pure product crystallized from the concentrated solution. Yield: 13.4 g (100%). ¹H NMR: (CDCl₃, 200 MHz) δ 7.75-7.15 (8H, m, ArH), 4.55-4.05 (5H, m, ArCHCH₂, H2, H4), 3.65-3.45 (2H, m, 2 H5), 2.35-2.10 (2H, m, 2 H3).

EXAMPLE 2

N-Fmoc-3-Hydroxypyrrolidine-5-Methanol

To a solution of N-Fmoc-trans-4-hydroxy-L-proline (13.4 g, 38.1 mmol) in 250 ml THF was added borane-methyl sulfide (78 mmol, 5.78 g, 7.22 ml) dropwise at room temperature. After the evolution of H₂ had ceased, the solution was heated to reflux with mechanical stirring. After 1 hour a white precipitate had formed. Methanol was carefully added, and the resulting solution refluxed for a further 15 minutes. The solution was cooled to room temperature, the solvents evaporated under reduced pressure, and the residual gum coevaporated with 2x100 ml methanol. The resulting crystalline product weighed 12.0 g (35.3 mmol, 93%). ¹H NMR: (CDCl₃, 200 MHz) δ 7.85-7.25 (8H, m, ArH), 4.50-4.10 (5H, m, ArCHCH₂, H3, H5), 3.80-3.40 (4H, m, 2 H2, 2 H6), 2.15-1.95 (1H, m, H2a), 1.80-1.60 (1H, m, H2b).

EXAMPLE 3

N-Fmoc-5-Dimethoxytrityloxymethyl-3-Hydroxypyrrolidine

The diol, N-Fmoc-3-hydroxypyrrolidine-5-methanol (10.59g, 31.2 mmol) was coevaporated with dry pyridine (2 X 50 ml), redissolved in 200 ml dry pyridine, and cooled in an ice bath. Dimethoxytrityl chloride (11.0 g, 32.5 mmol) was

- 52 -

added in portions over 30 min, and the solution stirred at 0 °C overnight. Methanol was then added (10 ml), and the solvent removed under reduced pressure. The resulting gum was redissolved in toluene (100 ml), filtered to remove the pyridinium hydrochloride and taken to dryness again. The residue was dissolved in CH₂Cl₂ (300 ml), washed with 150 ml 0.1 M citric acid solution, 150 ml sat NaHCO₃, brine, and dried with MgSO₄ followed by evaporation. The residue was crystallized from methanol and dried to give (15.4 g, 23.9 mmol, 77%).

EXAMPLE 4**5-Dimethoxytrityloxymethyl-3-Hydroxypyrrolidine**

To a solution N-Fmoc-5-dimethoxytrityloxymethyl-3-hydroxypyrrolidine (3.40 g, 5.30 mmol) in 15 ml DMF was added piperidine (1.09 ml, 0.935 g, 11.0 mmol). The solution was stirred at room temperature for 1 hour, water (100 ml) added, and the aqueous solution extracted with ethyl acetate (2 X 75 ml). The organic extracts were washed with aqueous NaHCO₃, brine, dried with MgSO₄ and evaporated. The residue was purified by flash column chromatography using a gradient of 1 - 3% MeOH in CH₂Cl₂ containing 0.5% triethylamine. Pure product was obtained (1.86 g, 84%). ¹H NMR: (CDCl₃, 200 MHz) δ 7.42-6.80 (13 H, ArH), 4.35 (1H, m, H5), 3.77 (6H, s, 2 OCH₃), 3.62 (1H, m, H3), 3.13-2.88 (4H, m, 2 H6, 2 H2), 1.87 (1H, q, H4a), 1.65 (1H, m, H4b).

EXAMPLE 5**N-(Phenylacetyl)-5-Dimethoxytrityloxymethyl-3-Hydroxypyrrolidine**

Phenylacetic acid (1.50 g, 11 mmol) and HOBt (1.63 g, 12 mmol) were dissolved in 100 ml CH₂Cl₂ and EDC (15 mmol, 2.88 g) was added. After 15 min, 5-dimethoxytrityloxymethyl-3-hydroxypyrrolidine was added, followed by DIEA (20 mmol, 3.5

ml). The reaction was stirred until the starting material was consumed, and quenched with 10 ml NaHCO_3 . The mixture was extracted twice with ethyl acetate, washed with NaHCO_3 , brine, dried with MgSO_4 , and evaporated. The product was purified by
5 flash chromatography to give 4.0 g product (75%). ^1H NMR: (CDCl_3 , 200 MHz) (2 rotamers, 3'-O-TMS) d 7.43-7.13, 6.88-6.74 (13 Ar-H), 4.67, 4.49, 4.37, 4.13 (4 m, 2H, H3, H5), 3.78 (s, 6H, OCH_3), 3.78-3.50 (m, 2H, H2a, b), 3.66, (s, 2H, CH_2Ar) 3.35 (q, 1H, H6a), 3.12 (m, 1H, H6b), 2.14-1.70 (m, 2H, H4a, 10 b), 0.10 (d, 9H, $\text{OSi}(\text{CH}_3)_3$).

EXAMPLE 6

Succinic acid Fluorenylmethyl Ester

Fluorenemethanol (10.0 g, 51.0 mmol) was dissolved in 150 ml CH_2Cl_2 , and succinic anhydride (5.6 g, 56 mmol) was
15 added. The solution was stirred for 6 h, and a further portion of succinic anhydride (2.5 g, 25 mmol) was added, and stirring continued overnight. The reaction appeared complete by TLC. The solvent was then removed, and the residue extracted with ethyl acetate, washed with 1% HCl, water,
20 brine, dried (MgSO_4) and evaporated to an oil which crystallized on standing. A quantitative yield of crude product was obtained which was used without further purification.

EXAMPLE 7

(N1-Thymine)-2-Acetic Acid

Methyl bromoacetate (25.5 g, 15.2 ml, 160 mmol) was added to a suspension of K_2CO_3 (44.2 g, 320 mmol) and thymine (20.2 g, 160 mmol) in 500 ml dry DMF with stirring overnight. The suspension was filtered and the solvent removed under
30 reduced pressure. The residue was suspended in 120 ml H_2O and 30 ml 4 N HCl, stirred for 30 minutes and filtered again. The solid was suspended in 250 ml H_2O , to which was added 100

- 54 -

ml 2.5 M NaOH. The solution was heated to boiling, cooled and acidified to pH 1 with concentrated HCl. The precipitate was dried in vacuo to give 13.6 g (73.6 mmol, 46%) pure product. ¹H NMR: (DMSO-d₆, 200 MHz) δ 7.48 (s, 1H, H₆), 4.37 (s, 2H, CH₂), 1.76 (s, 3H, CH₃).

EXAMPLE 8**N-Fmoc-3-Aminopropionic Acid**

Sodium bicarbonate (2.52 g, 30 mmol) and 3-aminopropionic acid (1.00 g, 11.2 mmol) were dissolved in 50 ml water and 50 ml dioxane was added. A solution of fluorenylmethyl chloroformate (3.10 g, 12.0 mmol) in 50 ml dioxane was added dropwise with stirring. After 6 hours the solution was diluted with water (100 ml) and saturated bicarbonate solution (50 ml), extracted once with diethyl ether, and the aqueous layer acidified to pH 2 with concentrated HCl. The cloudy solution was extracted with ethyl acetate (2 X 100 ml), washed with brine and dried with MgSO₄. After evaporation a mixture of the title product and the peptide dimer was obtained. The pure product was obtained by flash chromatography. ¹H NMR: (CDCl₃, 200 MHz) δ 7.95-7.26 (8H, m, ArH), 7.40-7.15 (3H, m, CHCH₂O), 3.20 (2H, t, J=8 Hz, CH₂N), 2.40 (2H, t, J=8 Hz, HOOCCH₂).

EXAMPLE 9**N-Imidazolyl-2-Acetic acid**

Imidazole (3.7 g, 54 mmol) was added to a suspension of sodium hydride (2.6 g of a 60% dispersion in oil, 60 mmol) in 50 ml dry THF. Bromoacetic acid (3.4 g, 24 mmol) was then added and the mixture stirred overnight. Water (1 ml) was then added and the solvent removed under reduced pressure. The residue was taken up in water (50 ml, pH >10), extracted with ether and the organic layer discarded. The aqueous layer was acidified to pH 1 with concentrated HCl and

- 55 -

extracted again with ether. The aqueous layer was evaporated to dryness. The oily residue was dissolved in absolute ethanol (EtOH) to precipitate NaCl, and recrystallized from acetone/methanol to give 1.22 g (7.5 mmol, 30%) pure product as the hydrochloride. ¹H NMR: (DMSO-d₆, 200 MHz) δ 9.20 (s, H₂), 7.76 (d, J = 1.5 Hz), 7.69 (d, J = 1.5 Hz), 5.20 (s, CH₂).

EXAMPLE 10**(9-Adenine)-2-Acetic Acid Ethyl Ester**

Sodium hydride (8.20 g 60% in oil, 205 mmol) was added to a suspension of adenine (25.0 g, 185 mmol) in 500 ml DMF. After vigorous stirring for 2 hours using a mechanical stirrer, H₂ evolution stopped and a thick slurry was obtained. Ethyl bromoacetate (55.6 g, 36.9 ml, 333 mmol) was added dropwise over 3 hours, and stirring continued for a further 1 hour. Water (10 ml) and H₂SO₄ were added to pH 4. The solvent was evaporated and the residue suspended in 500 ml H₂O, filtered and washed with water. The residue was recrystallized from 400 ml ethanol to give 23.8 g (108 mmol, 58%) pure product.

EXAMPLE 11**(N6-Benzoyl-9-Adenine)-2-Acetic Acid**

To a suspension of (9-adenylyl)-2-acetic acid ethyl ester (6.06 g, 27.4 mmol) in 250 ml dry pyridine was added benzoyl chloride (9.60 ml, 11.6 g, 82 mmol), and the solution stirred for 4 hours at room temperature. Methanol (25 ml) was added and the solvents evaporated. The residue was dissolved in ethyl acetate (2 X 250 ml), washed with 0.1 N HCl, H₂O, saturated NaHCO₃, brine, and dried with Na₂SO₄. The organic extracts were evaporated and the solid residue was redissolved in 250 ml THF at 0°C, to which was added 100 ml 1M NaOH. The solution was stirred at 0°C for 1 hour and

acidified to pH 1 with concentrated HCl, and the aqueous portion extracted once with ether. The product, which began to crystallize almost immediately, was collected by filtration to yield 4.96 g (61%). ¹H NMR: (DMSO-d₆, 200 MHz)
5 δ 8.86, 8.84 (d, H₂, H₈), 8.1 (d, 2H, J = 7.0 Hz, ArH), 7.69-7.58 (m, 3H, Ar-H), 5.22 (s, 2H, CH₂).

EXAMPLE 12

N-4-Benzoylcytosine

Cytosine hemihydrate (12.0 g, 100 mmol) was coevaporated
10 with pyridine and resuspended in 250 ml dry pyridine. Benzoyl chloride (58 ml, 70.3 g, 500 mmol) was added dropwise (exothermic). The solution was stirred at RT overnight, and water (50 ml) carefully added. The solvent was evaporated, and the residue dissolved in 700 ml H₂O containing 55 g NaOH.
15 The solution was stirred for 1 h after complete dissolution of the material. Concentrated HCl was then added to pH 4.0, the white precipitate was collected and boiled in 1 liter EtOH, cooled to RT and filtered to give 16.1 g product (75%).

EXAMPLE 13

20 N-4-Benzoyl-1-Cytosine-2-Acetic acid

To a suspension of N-4-Benzoylcytosine (15.0 g, 69.7 mmol) and K₂CO₃ (9.7 g, 70 mmol) in 500 ml DMF was added methyl bromoacetate (6.6 ml, 10.7 g, 70 mmol). The suspension was stirred vigorously for 3 days, filtered and
25 evaporated. The residue was treated with water (120 ml), and 10 ml 4N HCl for 15 min, and the solid collected by filtration. The residue was resuspended in 120 ml water, and 60 ml 2N NaOH added. The suspension was stirred at RT for 45 min, until all the solids had dissolved. The solution was
30 acidified to pH 2 with conc HCl, filtered, and the solid dried in vacuo at 60 °C to give 11.6 g product (61%).

EXAMPLE 14**N-2-Isobutyroyl-9-Guanine-2-Acetic Acid**

To a suspension of 2-amino-6-chloropurine (10 mmol) and K_2CO_3 (15 mmol) in DMF (25 ml) is added ethyl bromoacetate (10 mmol). The mixture is stirred vigorously for 24 hours, filtered and the solvent evaporated. The residue is resuspended in 25 ml pyridine and isobutyroyl chloride added (20 mmol). After stirring for 18 hours, water is added and the solvent removed. The residue is suspended in 1N HCl and heated to reflux for 1 hour. The suspension is then cooled to 0°C, NaOH added to pH 12, and the suspension stirred for 1 hour. The solution is acidified to pH 3, and the product is collected by filtration.

EXAMPLE 15**Benzyl 3,6,9,12-Tetraoxatridecanoate**

Triethyleneglycol monomethyl ether (10 mmol) and benzyl bromoacetate (11 mmol) are added to a suspension of anhydrous K_2CO_3 (15 mmol) in 50 ml anhydrous DMF. The suspension is stirred at room temperature overnight. Water is added and the emulsion is extracted with ethyl acetate (3 X 200 ml), washed with water, brine, and dried with $MgSO_4$. The solvent is evaporated and the residual oil purified by flash chromatography to give the title compound.

EXAMPLE 16**3,6,9,12-Tetraoxatridecanoic Acid**

Benzyl-3,6,9,12-Tetraoxatridecanoate (5 mmol) is dissolved in methanol (50 ml) and 10% palladium on carbon is added (100 mg catalyst/mmol). The suspension is shaken under 30 psi H_2 until the starting material is consumed. The suspension is filtered through a short pad of Celite, washed thoroughly with methanol, and the solvent evaporated. The product is used directly without purification.

EXAMPLE 17**Benzyl Bis-[(2-Pyridyl)-2-ethyl]-Aminoacetate**

To a suspension of K_2CO_3 (15 mmol) in 25 ml DMF was added 2,2'-bis(2-pyridylethyl)-amine (10 mmol) followed by benzyl bromoacetate (12 mmol). The suspension was stirred for 4 hours at room temperature. Water was then added, and the suspension extracted with ethyl acetate (2 X 100 ml), washed with 5% Na_2CO_3 , water, brine, dried with $MgSO_4$, and the solvents removed. The product was obtained as an oil in quantitative yield. Product was identified by NMR.

EXAMPLE 18**Bis(2-(2-Pyridyl)ethyl)-Aminoacetic Acid**

Benzyl bis-[(2-pyridyl)-2-ethyl]-aminoacetate (5 mmol) is dissolved in methanol (50 ml) and 10% palladium on carbon is added (100 mg catalyst/mmol). The suspension is shaken under 30 psi H_2 until the starting material is consumed. The suspension is filtered through a short pad of Celite, washed thoroughly with methanol, and the solvent evaporated. The product is used directly without purification.

EXAMPLE 19**N-Carbazolyl-2-Acetic acid**

The title compound is prepared as per Example 13 using carbazole as the starting heterocycle.

EXAMPLE 20**N-Pyrrolyl-2-Acetic acid**

The title compound is prepared as per Example 13 using pyrrole as the starting heterocycle.

EXAMPLE 21**N-Trifluoroacetyl-Glycine Triethylammonium salt**

To a suspension of glycine (1.50 g, 20 mmol) in 100 ml

- 59 -

dry methanol were added triethylamine (3.5 ml, 2.5 g, 25 mmol) and ethyl trifluoroacetate (3.0 ml, 3.55g, 25 mmol). The mixture was stirred overnight to give a homogeneous solution. The solvents were removed and the resulting oil
5 coevaporated with toluene to remove traces of methanol. The product was used without purification.

EXAMPLE 22**2-O-(Dimethoxytrityl)ethanol**

A solution of ethylene glycol (2.45 ml, 44 mmol) in dry
10 pyridine (25 ml) was cooled to 0 °C in an ice bath. Excess triethylamine (7 ml) and 4-dimethylaminopyridine catalyst (120 mg, 1 mmol) was added followed by the slow addition of dimethoxytrityl chloride (7.42 g, 21.9 mmol) over 30 minutes. The mixture was stirred at 0 °C for 1 hr and then room
15 temperature for 1 hr. The resulting solution was quenched with methanol and evaporated to dryness under reduced pressure. The residue was dissolved in saturated NaHCO₃ and extracted with EtOAc. The EtOAc extracts were washed with cold saturated sodium bicarbonate and brine. The organic
20 phase was separated, dried over sodium sulfate, filtered and evaporated under reduced pressure. The resulting residue was purified by flash column chromatography on silica gel using ethyl acetate-hexanes (gradient 10 to 20%). The title compound was isolated to yield 5.53 g (70%). ¹H NMR: (CDCl₃)
25 δ 7.50-7.20, 6.90-6.80 (m, 13 H, ArH), 3.80 (s, 6 H, OCH₃), 3.75 (t, 2H, CH₂OH), 3.25 (t, 2H, DMTOCH₂).

EXAMPLE 23**2-Dimethoxytrityl ethanol hemisuccinate Triethylammonium salt**

A solution of 2-O-(dimethoxytrityl)ethanol (1.0 g, 2.77
30 mmol), triethylamine (0.4 ml, 3 mmol), and 4-dimethylaminopyridine catalyst (120 mg, 1 mmol) in dry dichloroethane was treated with succinic anhydride (410 mg,

- 60 -

0.41 mmol). The mixture was stirred at room temperature for 16 hrs. The mixture was filtered and the filtrate was purified by silica gel flash column chromatography using chloroform-methanol-triethylamine to yield the title compound as a triethylammonium salt. ¹H NMR: (CDCl₃) δ 7.50-7.20, 6.90-6.80 (m, 13 H, ArH), 4.26 (t, 2H, CH₂OCO), 3.80 (s, 6 H, OCH₃), 3.25 (t, 2H, DMTOCH₂), 3.05 (q, 6H, N(CH₂CH₃)₃), 2.70 (m, 4H, OOCCH₂CH₂COO), 1.25 (t, 9H, N(CH₂CH₃)₃).

EXAMPLE 24**2-O-(Dimethoxytrityl)ethoxyphosphonic Acid**

A solution of imidazole (4.29 g, 63 mmol) in dry acetonitrile at 0 °C (100 ml) was treated dropwise with PCl₃ (1.77 ml, 20.3 mmol) over a period of 30 minutes. The resulting solution was further treated with triethylamine (9.06 ml, 65 mmol). To the thick slurry was added 2-O-(dimethoxytrityl)ethanol (2.10 g, 5.81 mmol) in anhydrous acetonitrile (150 ml) slowly over a period of 30 minutes. The mixture was allowed to warm to room temperature and stirred for 15 minutes. The mixture was quenched with 1M TEAB and the mixture is evaporated in vacuo to a minimum volume and extracted with dichloromethane (2x150 ml). The dichloromethane extracts were washed with TEAB and evaporated in vacuo. The residue was purified by flash column chromatography using a gradient of 0% to 5% methanol in dichloromethane/1% triethylamine to yield 1.3 g purified material (43%). ¹H NMR: (CDCl₃) δ 7.50-7.20, 6.90-6.80 (m, 13 H, ArH), 6.96 (d, 1H, J_{PH} = 624 Hz, PH), 4.06 (m, 2H, CH₂OP), 3.80 (s, 6 H, OCH₃), 3.25 (t, 2H, DMTOCH₂), 3.05 (q, 6H, N(CH₂CH₃)₃), 1.25 (t, 9H, N(CH₂CH₃)₃). ³¹P NMR (CDCl₃); 5.89.

EXAMPLE 25**Synthesis of 2-O-(Dimethoxytrityl)-ethylsuccinate Half Ester**

A solution of 2-O-(dimethoxytrityl)ethanol (1.0 g, 2.77

- 61 -

mmol), triethylamine (0.4 ml, 3 mmol), and 4-dimethylaminopyridine catalyst (120 mg, 1 mmol) in dry dichloroethane was treated with succinic anhydride (410 mg, 0.41 mmol). The mixture was stirred at room temperature for 16 hrs. The mixture was filtered and the filtrate was purified by silica gel flash column chromatography using chloroform-methanol-triethylamine to yield the title compound as a triethylammonium salt. ¹H NMR: (CDCl₃) δ 7.50-7.20, 6.90-6.80 (m, 13 H, ArH), 4.26 (t, 2H, CH₂OCO), 3.80 (s, 6 H, OCH₃), 3.25 (t, 2H, DMTOCH₂), 3.05 (q, 6H, N(CH₂CH₃)₃), 2.70 (m, 4H, OOCCH₂CH₂COO), 1.25 (t, 9H, N(CH₂CH₃)₃).

EXAMPLE 26**Derivatization of LCAA CPG With 2-O-(Dimethoxytrityl)-ethylsuccinate Half Ester**

2-O-(Dimethoxytrityl)ethylsuccinate half ester triethylammonium salt (135 mg) was dissolved in dichloromethane (5 ml). 4-Dimethylaminopyridine catalyst (40 mg, 0.2 mmol) was added followed by toluene diisocyanate (.029 ml, 0.2 mmol). The mixture was shaken for 18 min. Long chain alkyl amine controlled pore glass (LCAA CPG) (1.0 g) was added and the mixture was shaken with the exclusion of light for 16 hrs. The mixture was filtered and washed with dichloromethane and then diethylether (3x10 ml each). The CPG was shaken for 16 hrs in pyridine/water (4:1), filtered, and rinsed with pyridine (5x5 ml). A 10 mg sample of the dried CPG was treated with 3% trichloroacetic acid in dichloromethane. The presence of the trityl ion qualitatively verified the derivatization. The loading was measured to be 30 μmol/g by measuring the absorbance of the dimethoxytrityl cation.

EXAMPLE 27**Synthesis of 10-O-(Dimethoxytrityl)-1-decanol**

A solution of decane-1,10-diol in dry pyridine and

- 62 -

containing excess triethylamine is treated with one equivalent of dimethoxytrityl chloride for a period of six hours. The resulting solution is evaporated to dryness under reduced pressure, the residue redissolved in methylene chloride and the solution washed with cold saturated sodium bicarbonate, water and brine. The organic phase is separated, dried over sodium sulfate, filtered and again evaporated under reduced pressure. The resulting residue is flash-chromatographed on silica gel using ethyl acetate-hexanes to isolate the purified product. Characterization by H-NMR yields signals for the DMT group (multiplet, 8.0-7.0 ppm), the decane group (multiplets, 1.2-4.0 ppm) and the alcohol (variable).

EXAMPLE 28**15 Synthesis of 10-O-(Dimethoxytrityl)decyloxy-phosphonic acid**

A solution of three equivalents of imidazole in dry acetonitrile is treated dropwise with one equivalent of PCl_3 over a period of 30 minutes. The resulting solution is further treated with excess triethylamine to drive the reaction to completion. After 1 hr the mixture is treated with a solution of one equivalent of 10-O-(dimethoxytrityl)decan-1,10-diol in dry acetonitrile and the mixture stirred at room temperature for an additional hour. This mixture is treated with an excess of a solution of triethylammonium bicarbonate, pH 8, to yield the title compound. The compound is purified by repeated extraction of the bicarbonate solution with ethyl acetate. Pooling and drying of the extracts over sodium bicarbonate followed by evaporation of the solvent under reduced pressure yields a compound which is used as such without further purification. Characterization by ^{31}P NMR (doublet, 6 ppm, $\text{JP-H} = 600$ Hz) and ^1H NMR yields signals for the DMT and the decane groups as for 10-O-(dimethoxytrityl)decandiol and signals for the

triethylammonium groups (doublet, triplet, 3.2-2.2 ppm).

EXAMPLE 29

Synthesis of 10-O-(Dimethoxytrityl)decylsuccinate Half Ester

5 A solution of 10-O-(dimethoxytrityl)decan-1,10-diol in dry dichloromethane is treated with one equivalent of succinic anhydride, excess triethylamine and 5 mole % of 4-dimethylaminopyridine catalyst. The mixture is stirred overnight under anhydrous conditions and then further diluted
10 with dichloromethane. This solution is washed with cold, saturated sodium bicarbonate, water and brine. The solution is then dried over magnesium sulfate, filtered and evaporated to dryness under reduced pressure. The resulting solid is purified by silica gel flash column chromatography using
15 ethyl acetate-methanol-triethylamine to yield the title compound as the triethylammonium salt. The free acid is obtained by repeated coevaporation of this material with wet methanol. Characterization by ¹H NMR yields signals for the DMT and decylene groups as for 10-O-(dimethoxytrityl)decan-
20 1,10-diol and signals for the succinic group (two closely spaced doublet of doublets, 2.5-3.0 ppm).

EXAMPLE 30

Derivatization of LCAA CPG with 10-O-(Dimethoxytrityl)decylsuccinate Half Ester

25 A commercially obtained sample of controlled pore glass derivatized with long chain alkylamine groups (LCAA CPG) is suspended in dry acetonitrile. In a separate dry container, 10-O-(dimethoxytrityl)decylsuccinate half ester is treated with two equivalents of pentafluorophenol, excess
30 triethylamine and two equivalents of dicyclohexyl carbodiimide. The mixture containing activated 10-O-(dimethoxytrityl)decylsuccinate half ester is stirred under argon for one hour and then added to the suspension of CPG

while maintaining anhydrous conditions. The mixture is then shaken gently for 6 hr, the supernatant is separated and the process is repeated twice more. The quantity of 10-O-(dimethoxytrityl)decylsuccinate half ester which is used in each treatment is based on the concentration of available amine groups per gram of LCAA CPG, generally found to be 25-40 mmol/gram. The CPG is then treated with a dilute solution of acetic anhydride in pyridine for 1 hr to cap all unreacted amine functionalities and then washed several times with acetonitrile. The extent to which this CPG has been derivatized is determined by treating an accurately weighed sample of the resulting CPG with 2% dichloroacetic acid in acetonitrile and measuring the absorbance of an aliquot of the supernatant at 498 nm.

15 EXAMPLE 31

2-N-Fmoc-2-Amino-1,3-Propanediol

2-Amino-1,3-propanediol (3.48g, 38.2 mmol) and NaHCO_3 (8.00 g, 95.2 mmol) are suspended in 150 ml H_2O /Dioxane (1:1). Fluorenylmethyl chloroformate (11.4g, 44.0 mmol) in 25 ml toluene is added dropwise. The temperature of the reaction is maintained below 25 °C during the addition. The mixture is stirred vigorously overnight, and then quenched with 50 ml saturated NaHCO_3 solution and 50 ml water. The solution is extracted with 100 ml diethyl ether. The aqueous layer is acidified to pH 1 with concentrated HCl , and extracted twice with ethyl acetate, and the organic extracts are washed with brine. The solution is dried with MgSO_4 , filtered and the solvent removed in vacuo. The crude material is purified by silica gel column chromatography to give the title compound.

30 EXAMPLE 32

1-O-Dimethoxytrityl-N-Fmoc-2-Aminopropan-3-ol

A solution of N-Fmoc-2-amino-1,3-propanediol (13.79g, 44

- 65 -

mmol) in dry pyridine (250 ml) is cooled to 0°C in an ice bath. Excess triethylamine (7 ml) and 4-dimethylamino-pyridine catalyst (120 mg, 1 mmol) is added followed by the slow addition of dimethoxytrityl chloride (14.8g, 44 mmol) over 30 minutes. The mixture is stirred at 0°C until complete. The resulting solution is quenched with methanol and evaporated to dryness under reduced pressure. The residue is dissolved in saturated NaHCO₃ and extracted with EtOAc. The EtOAc extracts are washed with cold saturated sodium bicarbonate and brine. The organic phase is separated, dried over sodium sulfate, filtered and evaporated under reduced pressure. The resulting residue is purified by flash column chromatography on silica gel to give the title compound.

15 EXAMPLE 33

1-O-Dimethoxytrityl-N-Fmoc-2-Aminopropane-3-O-phosphonic Acid

A solution of imidazole (4.29g, 63 mmol) in dry acetonitrile at 0 °C (300 ml) is treated dropwise with PCl₃ (1.77 ml, 20.3 mmol) over a period of 30 minutes. The resulting solution is further treated with triethylamine (9.06 ml, 65 mmol). To the thick slurry was added 1-O-Dimethoxytrityl-N-Fmoc-2-Aminopropan-3-ol (3.58g, 5.81 mmol) in anhydrous acetonitrile (150 ml) slowly over a period of 30 minutes. The mixture is allowed to warm to room temperature and stirred for 15 minutes. The mixture is quenched with pyridine/water 9:1 (100 mL) and the mixture is evaporated in vacuo to a minimum volume and extracted with dichloromethane (2x150 ml). The dichloromethane extracts are washed with water and evaporated in vacuo. The residue is purified by silica gel column chromatography using dichloromethane/MeOH/pyridine to give the title compound.

EXAMPLE 34

- 66 -

1-O-Dimethoxytrityl-N-Fmoc-2-Aminopropane-3-O-Succinate

1-O-Dimethoxytrityl-N-Fmoc-2-aminopropan-3-ol is treated with succinic anhydride as per the general procedure of Example 60 to give the title compound.

5 EXAMPLE 35**Derivatization of LCAA CPG With 1-O-Dimethoxytrityl-N-Fmoc-2-Aminopropane-3-O-Succinate**

1-O-Dimethoxytrityl-N-Fmoc-2-Aminopropane-3-O-Succinate ester is coupled onto LCAA CPG as per the general procedure
10 of Example 61 to give the derivatized resin.

EXAMPLE 36**1-O-Dimethoxytrityl-2-Aminopropan-3-ol Succinate Derivatized Resin**

The Fmoc protecting group on the 2-amino group of the 1-
15 O-dimethoxytrityl-N-Fmoc-2-amino-1,3-propan-3-ol succinate derivatized CPG is removed by treatment with piperidine in dimethylformamide (DMF). CPG bound 1-O-dimethoxytrityl-N-Fmoc-2-amino-1,3-propanediol is treated with 2 equivalents of piperidine in DMF. The CPG is then washed with acetonitrile/
20 pyridine 1:1 and then treated a second time with 2 equivalents of piperidine in DMF. Finally, the CPG is washed with acetonitrile-pyridine and then acetonitrile to give the deprotected material.

EXAMPLE 37**25 1-O-Dimethoxytrityl-2-N-(acetylthymine) amino-1,3-Propanediol Succinate Derivatized Resin****Method A**

The 1-O-dimethoxytrityl-2-aminopropane-3-ol succinate derivatized resin (2.0 g, 1.0 mmol/gm loading, 1% cross-
30 linked) is swollen in dichloromethane (200 mL) and to this is added thymine-2-acetic acid (2.0g, 10 mmol), [O-(7-azabenz-

- 67 -

triazol-1-yl)-1,1,3,-tetramethyluronium hexafluorophosphate (3.8g, 10 mmol) and triethylamine (2.0g, 20 mmol). After 30 to 60 minutes the resin is washed 5 times with dichloromethane (50 mL), then 3 times with diethyl ether (100 mL), and is dried under a stream of nitrogen. The free flowing resin powder is used as is.

Method B

The 1-O-dimethoxytrityl-2-amino-1,3-propanediol succinate derivatized resin (2.0 g, 1.0 mmol/gm loading, 1% cross-linked) is swollen in dichloromethane (200 mL) and to this is added HOBt (0.1 M), PyBOP (0.1 M), N-methylmorpholine (0.15 M), as solutions in DMF followed by thymine-2-acetic acid (2.0g, 10 mmol). Coupling is allowed to proceed for 2-3 hours or overnight. The resin is washed 5 times with dichloromethane (50 mL), then 3 times with diethyl ether (100 mL), and is dried under a stream of nitrogen. The free flowing resin powder is used as is.

EXAMPLE 38

Sequential Addition and functionalization of n Backbone segments

The dimethoxytrityl protecting group of the derivatized resin of Example 37 is removed by a treatment with a solution of trichloroacetic acid (3% w/v) in dichloromethane. The solution is passed over the solid support until the DMT cation color is completely gone. The solid support is washed with dichloromethane until no trace of acid remains. The resin is then washed with acetonitrile-pyridine (4:1) followed by a simultaneous treatment of the CPG with 10 equivalents of 1-O-dimethoxytrityl-N-Fmoc-2-amino-3-O-phosphonic acid and 30 equivalents of adamantane carbonyl chloride in acetonitrile-pyridine (1:1). The mixture is agitated by circulating the reagents in the synthesis vessel for 2 minutes. The CPG is then briefly washed with

- 68 -

acetonitrile-pyridine and then treated with diisopropyl phosphite adamantane carbonyl chloride to cap all unreacted hydroxyl groups. The CPG is washed with acetonitrile-pyridine and then acetonitrile. The resulting phosphonic acid diester is reacted with a large molar excess of 10% diethylamine (the amine letter) in carbon tetrachloride/pyridine (1:1). The solid support is shaken for 15 minutes and the supernatant is removed by filtration. The solid support is washed with pyridine. A second treatment with diethylamine in carbon tetrachloride/ pyridine followed by shaking will ensure efficient oxidation to the phosphoramidate. The Fmoc protecting group is removed as per the general procedure of Example 62. The resulting free amine group is treated with thymine-2-acetic acid as per the procedure of Example 37. This procedure is repeated twice to give a dimer having thymineacetyl groups corresponding to the letter and the tether covalently bound to the amine group attached to carbon in the backbone segment. The functional groups bound to the phosphoramidate nitrogen are ethyl groups. This procedure when repeated n times will give a fully functionalized oligomer that is n+1 backbone segments long.

Upon completion of the addition of the last of the desired length and configuration of oligomeric sequence, the solid support is washed with pyridine/acetonitrile and the phosphoramidate is cleaved from the resin by treatment with concentrated ammonium hydroxide at room temperature for 3 hours. Evaporation of the supernatant and purification of the phosphoramidate on an RP-18 HPLC column yields the final oligomer.

EXAMPLE 39

N-Fmoc-Aspartic Acid- β -Benzyl Ester

Aspartic acid- β -benzyl ester (150 mmol) and diisopropyl-

- 69 -

ethylamine (66.3 ml, 49.1 g, 380 mmol) are suspended in 150 ml H₂O + 300 ml dioxane. Fluorenylmethyl chloroformate (43.25 g, 1.1 eq) in 100 ml dioxane is added dropwise. The temperature of the reaction is not allowed to rise above 10 °C during the addition. The mixture is stirred vigorously overnight, and most of the solvent removed in vacuo. Water and satd bicarbonate solution are added (250 ml each), and the solution extracted with 250 ml diethyl ether, which is discarded. The aqueous layer is acidified to pH 1 with conc HCl, and extracted twice with ethyl acetate (2 X 300 ml), and the organic extracts washed with brine. The solution is dried with MgSO₄, filtered and the solvent removed in vacuo to give the title compound.

EXAMPLE 40**15 4-Hydroxy-2-N-Fmoc-aminobutanoic acid**

2-N-Fmoc-aspartic acid- β -benzyl ester (10 mmol) is dissolved in dry THF (100 ml), cooled to 0 °C and Lithium borohydride (15 mmol) added. The solution is stirred at 0 °C and then room temperature until the complete disappearance of the starting material. Excess ethyl acetate is then added, and the solution is washed with 0.1M citric acid solution, brine and dried with MgSO₄. The crude material is purified by flash chromatography to give the title compound.

EXAMPLE 41**25 4-O-Dimethoxytrityl-2-N-Fmoc-aminobutanoic acid**

4-Hydroxy-2-N-Fmoc-aminobutanoic acid (30 mmol) is coevaporated with dry pyridine (2 X 50 ml), redissolved in 200 ml dry pyridine, and cooled in an ice bath. Dimethoxytrityl chloride (22.0 g, 65 mmol) is added in portions over 30 min, and the solution stirred at RT overnight. Water is then added (10 ml), and the solution stirred until the trityl ester is completely hydrolyzed. The

- 70 -

solvent is removed under reduced pressure. The residue is dissolved in CH_2Cl_2 (300 ml), washed with 150 ml 0.1 M citric acid solution, 150 ml sat NaHCO_3 , brine, and dried with MgSO_4 , followed by evaporation. The residue is purified by flash chromatography.

EXAMPLE 42**4-O-Dimethoxytrityl-2-N-Fmoc-aminobutan-1-ol**

To a stirred solution of 4-O-Dimethoxytrityl-2-N-Fmoc-aminobutanoic acid (140 mmol) in 500 ml THF is added Borane-methyl sulfide (290 mmol, 21.8 g, 27.3 ml) dropwise at RT. Stirring is continued until the reaction is complete. Methanol is carefully added (vigorous H_2 evolution), and the resulting solution stirred for a further 15 min. The solvent is evaporated under reduced pressure, and the residual gum coevaporated with 2 X 300 ml MeOH. The product is purified by flash chromatography.

EXAMPLE 43**1-O-Dimethoxytrityl-2-N-Fmoc-2-Aminobutane-4-O-Phosphonic Acid**

1-O-Dimethoxytrityl-2-N-Fmoc-2-aminobutan-4-ol is treated as per the general procedure of Example 59 to give the title compound.

EXAMPLE 44**1-O-Dimethoxytrityl-2-N-Fmoc-2-Aminobutan-4-O-Succinate.**

1-O-Dimethoxytrityl-2-N-Fmoc-2-aminobutan-1-ol is treated as per the general procedure of Example 60 to give the title compound.

EXAMPLE 45

1-Derivatization of LCAA CPG With 1-O-Dimethoxytrityl-2-N-Fmoc-2-Aminobutane-4-O-Succinate

- 71 -

1-O-Dimethoxytrityl-2-N-Fmoc-2-aminobutane-4-O-succinate is treated as per the general procedure of Example 61 to give the derivatized resin.

EXAMPLE 46

5 **1-O-Dimethoxytrityl-2-Aminobutane-4-O-Succinate Derivatized Resin**

The derivatized resin of Example 45 is treated as per the general procedure of Example 62 to remove the Fmoc protecting group giving the title compound attached to resin.

10 **EXAMPLE 47**

1-O-Dimethoxytrityl-2- (N-2-Acetylthymine) -Aminobutane-4-O-succinate Derivatized Resin

15 The derivatized resin of Example 46 is treated with N-1-thymine-2-acetic acid as per the procedure of Example 37 to give the title compound attached to resin.

EXAMPLE 48

Synthesis of a 3-mer Having the 2-Amino-1,4-Butanediol backbone segment

20 1-O-Dimethoxytrityl-2- (N-2-acetylthymine)-aminobutane derivatized resin is treated with 1-O-dimethoxytrityl-2-N-Fmoc-2-amino-4-phosphonic acid-1,4-butanediol, morpholine, and N-1-thymine acetic acid as per the procedures of Example 38 and Example 126 to give a 3 mer with 2-N-acetylthymine bound to the amino groups and morpholine groups as the
25 phosphoramidate substituent.

EXAMPLE 49

2- (N-Fmoc) -Glutamic acid-γ-methyl ester

30 Glutamic acid-γ-methyl ester (150 mmol) and diisopropylethylamine (66.3 ml, 49.1 g, 380 mmol) are suspended in 150 ml H₂O + 300 ml dioxane. Fluorenylmethyl chloroformate (

- 72 -

43.25 g, 1.1 eq) in 100 ml dioxane is added dropwise. The temperature of the reaction is not allowed to rise above 10 °C during the addition. The mixture is stirred vigorously overnight, and most of the solvent removed *in vacuo*. Water and satd bicarbonate solution are added (250 ml each), and the solution extracted with 250 ml diethyl ether, which is discarded. The aqueous layer is acidified to pH 1 with conc HCl, and extracted twice with ethyl acetate (2 X 300 ml), and the organic extracts washed with brine. The solution is dried with MgSO₄, filtered and the solvent removed in *vacuo* to give the title compound.

EXAMPLE 50**5-Hydroxy-4-N-Fmoc-aminopentanoic acid methyl ester**

To a solution of 2-(N-Fmoc)-glutamic acid-γ-methyl ester (140 mmol) in 500 ml THF is added Borane-methyl sulfide (290 mmol, 21.8 g, 27.3 ml) dropwise at RT (3 neck flask, mechanical stirrer, condenser, dropping funnel). After the evolution of H₂ has ceased, the solution is heated to reflux with vigorous stirring. After 1 hr a white precipitate has formed. Methanol is carefully added (vigorous H₂ evolution), and the resulting solution refluxed for a further 15 min. The solution is cooled to RT, the solvents evaporated under reduced pressure, and the residual gum coevaporated with 2 X 300 ml MeOH. The product is purified by flash chromatography.

EXAMPLE 51**5-O-Dimethoxytrityl-4-Fmoc-aminopentanoic acid methyl ester**

5-Hydroxy-4-N-Fmoc-aminopentanoic acid methyl ester (30 mmol) is coevaporated with dry pyridine (2 X 50 ml), redissolved in 200 ml dry pyridine, and cooled in an ice bath. Dimethoxytrityl chloride (11.0 g, 32.5 mmol) is added in portions over 30 min, and the solution stirred at 0°C

- 73 -

overnight. Methanol is then added (10 ml), and the solvent removed under reduced pressure. The resulting gum is redissolved in toluene (100 ml), filtered to remove the pyridinium hydrochloride and taken to dryness again. The residue is dissolved in CH_2Cl_2 (300 ml), washed with 150 ml 0.1 M citric acid solution, 150 ml sat NaHCO_3 , brine, and dried with MgSO_4 followed by evaporation. The residue is purified by flash chromatography to give the title compound.

10 **EXAMPLE 52****5-O-Dimethoxytrityl-4-Fmoc-aminopentan-1-ol**

5-O-Dimethoxytrityl-4-Fmoc-aminopentanoic acid methyl ester (10 mmol) is dissolved in dry THF (100 ml), cooled to 0 °C and Lithium borohydride (10 mmol) added. The solution is stirred at 0 °C and then room temperature until the complete disappearance of the starting material. Excess ethyl acetate is then added, and the solution washed with 0.1M citric acid solution, sat NaHCO_3 , brine and dried with MgSO_4 . The product is purified by flash chromatography

20 **EXAMPLE 53****5-O-Dimethoxytrityl-4-N-Fmoc-aminopentan-1-ol hydrogen phosphonate**

Imidazole (6.81 g, 100 mmol) is dissolved in 400 ml dry CH_3CN and cooled to 0 °C. Phosphorus trichloride (2.62 ml, 4.12 g, 30 mmol) is added dropwise, followed by triethylamine (21 ml, 15.2 g, 150 mmol). A thick slurry develops to which is added over 15 min a solution of 5-O-dimethoxytrityl-4-N-Fmoc-2-amino-1,5-pentanediol (10 mmol) in 50 ml CH_3CN . Once the addition is complete, the ice bath is removed and the solution stirred at RT for 30 min. The reaction is stopped by the addition of 100 ml pyridine/water (9:1). The solvent is removed and the residue extracted (3X200 ml) with CH_2Cl_2 , and washed with water. The organic phase is dried with MgSO_4

- 74 -

and concentrated under reduced pressure. The product is further purified by flash chromatography using a gradient of MeOH (1-10%) in CH_2Cl_2 + 1% pyridine.

EXAMPLE 54

5 **5-O-Dimethoxytrityl-4-N-Fmoc-Aminopentane-1-O-Succinic Acid Half Ester**

5-O-Dimethoxytrityl-4-N-Fmoc-aminopentane-1-ol is treated as per the general procedure of Example 60 to give the title compound.

10 **EXAMPLE 55**

Derivatization of LCAA CPG With 5-O-Dimethoxytrityl-4-N-Fmoc-Aminopentane-1-O-Succinic Acid Half Ester

5-O-Dimethoxytrityl-4-N-Fmoc-Aminopentane-1-O-Succinic Acid Half Ester is treated as per the general procedure of
15 Example 61 to give the derivatized resin.

EXAMPLE 56

5-O-Dimethoxytrityl-4-Aminopentane-1-O-Succinic acid Derivatized Resin

The derivatized resin of Example 55 is treated as per
20 the procedure of Example 62 to remove the Fmoc protecting group giving the title compound attached to resin.

EXAMPLE 57

5-O-Dimethoxytrityl-4-N-Phenylacetyl-Aminopentane-1-O-Succinic acid Derivatized Resin

25 The derivatized resin of Example 56 is treated with phenyl acetic acid as per the procedure of Example 38 to give the title compound attached to resin.

EXAMPLE 58

Synthesis of a 3-mer having the 2-amino-1,5-pentanediol

- 75 -

backbone segment

1-O-Dimethoxytrityl-2(phenylacetyl)-Amino-5-Phosphonic
Acid-1,5-pentanediol Derivatized Resin is treated with 1-O-
dimethoxytrityl-2-N-Fmoc-2-amino-5-phosphonic acid-1,5-
5 pentanediol, morpholine, and phenylacetic acid as per the
procedure of Example 38 and Example 126 to give a 3 mer with
phenylacetyl bound to the amino groups and morpholine groups
as the phosphoramidate substituent.

EXAMPLE 59

10 **General Procedure for Converting Aminodiol Monomer Subunits
to the H-Phosphonate Monoesters**

**N-Fmoc-5-Dimethoxytrityloxymethylpyrrolidine-3-O-
Hydrogen Phosphonate Triethylammonium salt**

Imidazole (6.81 g, 100 mmol) was dissolved in 400 ml dry
15 CH₃CN and cooled to 0 °C. Phosphorus trichloride (2.62 ml,
4.12 g, 30 mmol) was added dropwise, followed by
triethylamine (21 ml, 15.2 g, 150 mmol). A thick slurry
developed to which was added over 15 min a solution of N-
Fmoc-5-dimethoxytrityloxymethyl-3-hydroxypyrrolidine (10
20 mmol) in 50 ml CH₃CN. Once the addition was complete, the
ice bath was removed and the solution stirred at RT for 30
min. The reaction was stopped by the addition of 100 ml
Pyridine and 10 ml water. The solvent was removed and the
residue coevaporated with 100 ml Pyridine, then 100 ml
25 toluene. The residue was then dissolved in 0.05M TEAB,
extracted (3X200 ml) with CH₂Cl₂, and the extract washed with
0.05M TEAB. The organic phase was dried with MgSO₄ and
concentrated under reduced pressure. The product was further
purified by flash chromatography using a gradient of MeOH (1-
30 10%) in CH₂Cl₂ + 0.5 % TEA.

EXAMPLE 60

General Procedure for converting Aminodiol Monomer Subunits

- 76 -

to the Succinate derivatives

N-Fmoc-5-Dimethoxytrityloxymethylpyrrolidine-3-O-

Succinate

N-Fmoc-5-Dimethoxytrityloxymethylpyrrolidine (2.0 mmol),
5 succinic anhydride (300 mg, 3.0 mmol), DMAP (1.0 mmol, 120
mg) and triethylamine (0.4 ml, 3.0 mmol) are dissolved in
dichloromethane and stirred overnight. The solution is
extracted with dichloromethane, washed with 0.1M citric acid,
water, brine, dried and evaporated. The residue is filtered
10 through a short pad of silica and used directly.

EXAMPLE 61

**N-Fmoc-5-Dimethoxytrityloxymethylpyrrolidine-Controlled Pore
Glass (N-FMOC-5-DMT-hp-CPG)**

N-Fmoc-5-Dimethoxytrityloxymethylpyrrolidine-3-O-
15 Succinate is dissolved in dry dichloromethane (50 ml), DMAP
added (250 mg, 2 mmol) followed by toluenediisocyanate (288
ul, 2.0 mmol) . The mixture is swirled for 10 min then 10 g
LCAA-CPG is added followed by DIEA (2 mmol, 0,34 ml). The
suspension is kept in the dark and agitated periodically for
20 6-16 h. The solid is filtered, washed with dichloromethane
and ether, then suspended in 80 ml pyridine + 20 ml water.
After 1 h, the support is filtered, washed with dry pyridine
(5X), dichloromethane (3X) , and suspended in 60 ml
dichloromethane, to which 10 ml TEA, 10 ml acetic anhydride,
25 3 ml N-methylimidazole are added. After 1 h, the support is
filtered, washed extensively with dichloromethane and ether
and dried. The CPG is analyzed for loading by weighing a
portion of CPG , dissolving in 0.1M toluene sulfonic acid and
measuring the absorbance at 498 nm.

30 **EXAMPLE 62**

**Deprotection of Backbone Segment Amino Combinatorial Site
Protecting groups General Procedure**

- 77 -

The solid support (1-10 umol) bearing an Fmoc protected amine is washed with a 10 % (v/v) solution of piperidine in DMF for 15 seconds, and suspended in the piperidine solution for 15 minutes. The solvent is removed and a fresh portion of piperidine/DMF is added for a further 15 minutes. The solid support is then washed with several portions of DMF to remove all traces of piperidine.

Using this procedure the Fmoc protected backbone segment amino combinatorial site of the solid support bound N-Fmoc-5-dimethoxytrityloxymethylpyrrolidine of Example 61 is deprotected.

EXAMPLE 63

Coupling of Carboxylic acids to Backbone Segment Amino Combinatorial Sites General procedure

Deprotected backbone segment amino combinatorial sites are treated simultaneously with a solution of a carboxylic acid (0.4 mmol/ml) and diisopropylethylamine (0.8 mmol/ml) in DMF, and BOP reagent or HATU (0.4 mmol/ml) in DMF. A tenfold excess of reagents are added at the 10 umol scale level. The reaction is allowed to proceed for 30 minutes and a further ten equivalents of carboxylic acid and BOP is added. After 30 minutes, the solid support is washed with DMF until all the reagents are removed.

EXAMPLE 64

Removal of the DMT protecting group General procedure

Aminodiol monomeric subunits having dimethoxytrityl protecting group are deprotected using a solution of trichloroacetic acid (3%, w/v) in dichloromethane is passed over the solid support until the DMT cation color is completely gone. The solid support is washed with dichloromethane until no trace of acid remains.

- 78 -

EXAMPLE 65**Coupling of Sulfonyl Chlorides to Backbone Segment Amino
Combinatorial Site General procedure**

To the free amine on solid support obtained by
5 piperidine treatment of the Fmoc protected backbone segment
is added simultaneously a solution of a sulfonyl chloride
(0.1 mmol/ml) and diisopropylethylamine (0.25 mmol/ml) in
pyridine/CH₃CN. A tenfold excess of reagents is added at the
10 umol scale level. The reaction is allowed to proceed for
30 minutes, and the solid support was washed with
Pyridine/CH₃CN until all the reagents are removed.

EXAMPLE 66**Coupling of acyl groups to Backbone Segment Amino
Combinatorial Site General procedure**

15 To the free amine on solid support are added
simultaneously a solution of an activated acylating agent
(0.1 mmol/ml) and diisopropylethylamine (0.25 mmol/ml) in
Pyridine/CH₃CN. A tenfold excess of reagents are added at the
10 umol scale level. The reaction is allowed to proceed for
20 30 minutes, the solid support is washed with Pyridine/CH₃CN
until all the reagents are removed. Groups useful for
acylating the free amine combinatorial site include acid
halides, acid fluorides, acid imidazolides, acid anhydrides,
sulfonyl chlorides, chloroformates, isocyanates, isothio-
25 cyanates.

EXAMPLE 67**Phosphoramidate Library Synthesis General procedures**

The general procedures described below outline the
methods for the preparation of phosphoramidate combinatorial
30 libraries. All manipulations can be accomplished using an
automated synthesizer to deliver solvents and reagents to a
reaction vessel containing the reactants attached to a solid

- 79 -

support, usually controlled pore glass (CPG) or Tentagel (TG). The design of such synthesizers and the loading of the solid support allow the manipulations to be performed on scales ranging from 50 pmol to approximately 10 mmol if desired. The manipulations can also be carried out manually by using a syringe with a glass frit as the reaction vessel, by drawing the appropriate solutions into the syringe. The reactions described can be carried out on single compounds attached to the solid support, or on complex mixtures prepared by the technique of bead portioning/mixing. In brief, these techniques involve the addition of unique reagents to an equal number of separate portions of solid support. Once the individual reactions are complete, the portions of solid support are combined and mixed thoroughly as a slurry in an appropriate solvent and redivided into the number of portions required by the number of different reagents in the next step. In this way, a unique reagent is added to a mixture of compounds, to create all possible combinations of the different reagents.

Oxidation of H-Phosphonate Diester linkages To Form Phosphoramidate linkages having letters

Method A: Incorporation of Letters In Predetermined Sequence

The solid support (e.g. LCAA CPG) is derivatized with a first monomeric subunit or other group having a terminal reactive group capable of forming a covalent bond with a primary or secondary amine. Example 26 illustrates a non-aminodiol monomer subunit attached to a solid support and Example 45 illustrates attachment of an aminodiol monomer subunit to a solid support. When an aminodiol monomer subunit is derivatized to the solid support the amino group is deprotected using the general procedure of Example 62 and coupled with a letter having an optional tether as illustrated in Examples 36, 37, 63, and 65 (also see Examples

- 80 -

122-130). The hydroxyl protecting group is then removed as illustrated in Example 64 and the next desired aminodiol monomer subunit having a phosphonic acid monoester group on one of the hydroxyls is condensed onto the terminal free hydroxyl of the derivatized solid support as per the procedure of Example 48. The resulting phosphonic acid diester is reacted with a large molar excess of a primary or secondary amine in carbon tetrachloride/ pyridine to form the phosphoramidate having a letter attached thereto. The amine letter is added in a solution of carbon tetrachloride/pyridine. The solid support is washed with pyridine. A second treatment with a large molar excess of the amine letter in carbon tetrachloride/ pyridine followed by shaking will ensure efficient oxidation to the phosphoramidate. The amino protecting group on the second backbone substituent is removed using piperidine treatment and the free amino is coupled to a letter as above. In this case the procedures give an oligomeric compound two units in length.

Additional functionalized monomer units can be added by repeating the following steps as often as desired: 1) removal of the oxygen protecting group, 2) coupling of an additional scaffold phosphonate monoester, 3) oxidation of the phosphonate diester with carbon tetrachloride in the presence of the desired amine letter, 4) removal of the nitrogen protecting group on the last scaffold monomer unit, 5) functionalization of the amino group with the desired functionality.

The oxidative incorporation of letters to form phosphoramidate linkages can be performed stepwise or all at once for a uniform substitution of letters. For the addition of two adjacent like letters in the oligomeric structure, the oxidation step can be delayed until the backbone to support all of these letters is synthesized and all H phosphonate sites that will bear this letter are then oxidized simultaneously.

- 81 -

The above steps are repeated until all of the letters of the oligomer have been added. All letters are predetermined in this method of synthesis. Upon completion of the addition of the last of the desired length and configuration of the oligomeric compound, the solid support is washed with pyridine/acetonitrile and the phosphoramidate is cleaved from the resin by treatment with concentrated ammonium hydroxide at room temperature for 3 hours. Evaporation of the supernatant and purification of the phosphoramidate on an RP-18 HPLC column yields the final oligomer.

Method B: Incorporation of Phosphoramidate Letters in a Random Sequence

The method of oligomer synthesis as described above in Method A is repeated to synthesize the oligomer of desired length. To randomize the amine letters on the oligomer, the method of adding a letter as described in Method a above is followed except that, for randomization, the amine letters in carbon tetrachloride and a suitable cosolvent are added as a mixture, preferably one normalized for relative reactivity. Random distribution of amine letters from this mixture of amine letters is verified experimentally by treatment of an oligomer, which has been previously treated with a mixture of amine letters and subsequently worked up and purified, with 10% aqueous formic acid at 50-70 °C to release the amine letters. The actual percentages of incorporation of the individual amine letters is then determined by HPLC analysis of the reaction mixture and the relative individual rates are calculated. Having once determined the relative rates, in further iteration of the sequences, the concentration of amine letters within a mixture is adjusted to reflect these rate differences.

In a variation of this method of randomization, in a five mer all sites of which are to be randomized, the oxidation is effected simultaneously. The five mer backbone is synthesized as above and a mixture of the letters added.

- 82 -

Upon completion of the backbone synthesis, the oxidation of amine letters is effected on all five sites as a single step.

In a further variation of this method of randomization, upon completion of the synthesis of the first backbone segment, the resin is split into five portion and each portion is individually oxidized with one of the amine letter. The individual portions of the resin are recombined and a further backbone segment is attached thereby extending the oligomeric compound a further unit. The resin is then again split, and the individual portion each oxidized with one of the amine letter. This cycle is repeated to complete the synthesis.

Coupling of letters to the free amino positions is performed as in Method A for a predetermined sequence or Method B for a random sequence.

Method C: Incorporation of Amine Letters in Fixed/random Sequence

Combining methods A and B above can be used to fix certain positions while randomizing other positions as the oligomeric compound is synthesized. This method is further used in combination with a SURF™ combinatorial strategy.

The following is a representative list of amine letters:
allylamine

3,3-Dimethylaminopropylamine

1-Phenyl-1,3,8-Triazaspiro[4.5]decan-4-one

Azetidine

Benzylamine

Butylamine

L-(-)-2-Aminocaprolactam

Heptamethyleneimine

cyclopropylamine

Diethylamine

Dimethoxyethylamine

2,5-Dimethyl-N-phenylpiperazine

Methylaminomethyl-1,3-dioxolane

- 83 -

- N-Acetyl Ethylenediamine
3-Aminopropyl Imidazole
Methoxyethylamine
Morpholine
5 3-Aminopropyl Morpholine
Isonipecotamide
Piperonylamine
2-Aminomethylpyridine
Piperidine
10 Piperidone ethylene ketal
2-Aminoethyl-1-methylpyrrolidine
N(2-Pyridyl)piperazine
Piperonylpiperazine
3-Isopropoxypropylamine
15 Piperazine
Thienylethylamine
3-Trifluoromethoxybenzylamine
Tryptamine
Tetrahydrofurfurylamine

20 **EXAMPLE 68**

**1-Acetyl Thymine/Benzylamine Phosphoramidate Oligomer
Synthesis**

A solid support is derivatized with 2-O-(dimethoxytrityl)ethylsuccinate half ester as in Example 26. The DMT
25 protecting group is removed using the standard method of
Example 64. 4-O-Dimethoxytrityl-2-N-Fmoc-aminobutan-1-ol is
treated with PCl_3 as per Example 28 to form the phosphonic
acid which is condensed onto the derivatized resin as in
Example 38. The Fmoc amino protecting group is removed as
30 per Example 4 and (N1-thymine)-2-acetic acid (Example 7) is
coupled to the resulting free amino as per the method of
Example 38. The above methods of examples 64, 28, 37 and 38
are repeated until six of the above phosphonic acid residues

- 84 -

are incorporated. The resulting six mer is treated using the procedures of Example 67, with a large excess of benzylamine in carbon tetrachloride/pyridine. The solid support is shaken for 15 minutes and the supernatant is removed by
5 filtration and then washed with pyridine. A second treatment with a large excess of benzylamine in carbon tetrachloride/pyridine followed by shaking will insure efficient oxidation to the phosphoramidate. The resin is washed with
10 pyridine/acetonitrile and then the phosphoramidate is cleaved from the resin by a treatment with concentrated ammonium hydroxide at room temperature for 3 hours. Evaporation of the supernatant and purification of the phosphoramidate on an RP-18 HPLC column will yield the final oligomer. The
15 stepwise H phosphonate coupling efficiency is determined by measuring the absorbance of the trityl ion. The resulting six mer will have a benzylamine at each of the phosphoramidate linkages and will have an acetylthymine group at each of the backbone segment amino combinatorial sites.

EXAMPLE 69**20 Hydrogen phosphonate Coupling General Procedure**

A portion of solid support (CPG or other polymeric support e.g. TentaGel) derivatized with a DMT protected alcohol linked via a succinate linker (1 μ mol) is loaded into a DNA synthesis column, and attached to an automated DNA
25 synthesizer programmed to perform the following functions:
1) Wash with dichloromethane;
2) Treat with 3% trichloroacetic acid in dichloromethane to remove the DMT protecting group;
3) Wash with dichloromethane and CH₃CN/Pyridine (1:1);
30 4) Coupling: addition of alternating portions of 0.2 M Adamantoyl chloride or Pivaloyl chloride in CH₃CN/Py (1:1) and 0.05 M H-phosphonate monomer in CH₃CN/Py (1:1) for 1 min;
5) Wash with CH₃CN/Pyridine (1:1);
6) Stop and proceed to an oxidation procedure for oxidation

- 85 -

of each linkage independently or repeat steps 1 through 5 to add additional monomer subunits prior to oxidation.

The product of this sequence of reactions is an H-phosphonate diester, which is oxidized by one of several methods including those described below.

Oxidation procedure 1: Phosphodiester

The solid support-bound H-phosphonate diester is treated (manually or automatically) with equal volumes of solution A (0.2 M I_2 in THF) and solution B (N-methylmorpholine/ H_2O /THF 1:1:8) for 5 min, followed by equal volumes of solution A and solution C (TEA/ H_2O /THF 1:1:8) for 5 min, followed by washing with CH_3CN /Py (1:1).

Oxidation procedure 2: Phosphorothioate

The solid support-bound H-phosphonate diester is treated (manually or automatically) with a solution of S_8 in CS_2 /Lutidine for 30 min. The solid support is then washed with CH_3CN /Py (1:1).

Oxidation procedure 3: Phosphoramidate

The solid support-bound H-phosphonate diester is treated (manually or automatically) with a solution of the required amine (10 % V/V) in CCl_4 /Pyridine 1:1 for 15-30 min. The solid support is then washed with CH_3CN /Py (1:1).

EXAMPLE 70

Combinatorial library synthesis having phosphoramidate

linkages

N-Fmoc-5-dimethoxytrityloxymethyl-3-hydroxypyrrolidine (Example 2), 1-O-dimethoxytrityl-N-Fmoc-2-amino-1,3-propanediol (Example 32), and 4-O-dimethoxytrityl-2-N-Fmoc-aminobutan-1-ol (Example 42) are treated in separate reactions with succinic anhydride as per example 60 to give the corresponding succinyl derivatives. In a separate set of reactions each of the above aminodiol monomer subunits are

- 86 -

treated with PCl_3 as per Example 28 to give the corresponding H-phosphonate monoesters. A solid support is divided into three equal portions and each portion is treated with one of the succinyl derivatized aminodiol monomer subunits as per Example 61. The solid support is combined and the amino protecting groups are removed as per the general procedure of Example 62. The solid support is washed with pyridine/acetonitrile, dried, and redivided into three equal parts.

Each portion of the solid support is treated with one of (N1-Thymine)-2-acetic acid (Example 7), (N6-benzoyl-9-adenine)-2-acetic acid (Example 11), or N-4-benzoyl-1-cytosine-2-acetic acid (Example 13) following the general procedure of Example 63. The procedure of mixing, drying and redividing the solid support is repeated.

Each portion of solid support is treated with one of the H-phosphonate monoesters above as per the procedure of Example 30, to form the H-phosphonate diesters. The procedure of mixing, drying and redividing the solid support is repeated. Each portion of the solid support is treated with a large molar excess of one of benzylamine, 2-(2-aminoethyl)-1-methylpyrrolidine, or piperonyl amine in carbon tetrachloride/pyridine. The solid support is shaken for 15 minutes and the supernatant is removed by filtration and then washed with pyridine. A second treatment with a large excess of each amine letter in carbon tetrachloride/pyridine followed by shaking will insure efficient oxidation to the phosphoramidate. The solid support is combined and the amino protecting groups are removed as above. The solid support is washed with pyridine/acetonitrile, dried, and redivided.

Each portion of the solid support is treated with one of (N1-Thymine)-2-acetic acid (Example 7), (N6-benzoyl-9-adenine)-2-acetic acid (Example 11), or N-4-benzoyl-1-cytosine-2-acetic acid (Example 13) following the general procedure of Example 63. The procedure of mixing, deblocking the hydroxyl protecting group as per Example 64, drying and

- 87 -

redividing the solid support is repeated.

Each portion of solid support is treated with one of the H-phosphonate monoesters above as per the procedure of Example 69 to form the H-phosphonate diesters. The procedure of mixing, drying and redividing the solid support is repeated. Each portion of the solid support is treated with a large molar excess of one of benzylamine, 2-(2-aminoethyl)-1-methylpyrrolidine, or piperonyl amine in carbon tetrachloride/pyridine. The solid support is shaken for 15 minutes and the supernatant is removed by filtration and then washed with pyridine. A second treatment with a large excess of each amine letter in carbon tetrachloride/pyridine followed by shaking will insure efficient oxidation to the phosphoramidate. The solid support is combined and the amino protecting groups are removed as above. The solid support is washed with pyridine/acetonitrile, dried, and redivided.

Each portion of the solid support is treated with one of (N1-Thymine)-2-acetic acid (Example 7), (N6-benzoyl-9-adenine)-2-acetic acid (Example 11), or N-4-benzoyl-1-cytosine-2-acetic acid (Example 13) following the general procedure of Example 63.

The resulting oligomeric compounds are cleaved from the solid support by a treatment with concentrated ammonium hydroxide at room temperature for 3 hours. Evaporation of the supernatant and purification of the phosphoramidate linked oligomeric compounds on an RP-18 HPLC column will yield the final combinatorial libraries consisting of all the possible oligomeric compounds that can be prepared using the three aminodiols monomer subunits, the three functional groups, and the three amines.

EXAMPLE 71

Combinatorial library synthesis having phosphodiester or phosphorothioate linkages

- 88 -

A three mer is synthesized using the reagents and procedures of Example 68 except that oxidation of the H-phosphonate diester linkage is accomplished using the procedures illustrated in Example 69, Procedure 1 or 2, to give either uniform phosphodiester or uniform phosphorothioate linkages.

EXAMPLE 72

Combinatorial library synthesis having mixed phosphodiester, phosphorothioate, and phosphoramidate linkages

A three mer is synthesized using the reagents and procedures of Example 68 except that oxidation of the H-phosphonate diester linkage is accomplished by dividing the solid support into three equal portions and using the procedures illustrated in Example 69, Procedure 1, 2, or 3 to give phosphodiester, phosphorothioate, or phosphoramidate linkages.

EXAMPLE 73

Standard Oligomer Coupling Cycle Using Standard DNA Synthesis Protocols

The oligomeric compounds of the invention are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) as is done with standard oligonucleotides using standard phosphoramidate chemistry with oxidation by iodine (see, Oligonucleotide Synthesis, A Practical Approach, M. J. Gait., ed., Oxford University Press, New York, 1990). For phosphorothioate oligomeric compounds, the standard oxidation bottle is replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the step wise thiation of the phosphite linkages. The thiation wait step is increased to 68 sec and is followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55 °C (18 hours), the

oligomeric compounds can be purified by precipitation twice out of 0.5 M NaCl solution with 2.5 volumes ethanol or by HPLC chromatography using a RP-18 column. Analytical gel electrophoresis is effected in 20% acrylamide, 8 M urea, 454 mM Tris-borate buffer, pH = 7.0. Phosphodiester and phosphorothioate oligomeric compounds are judged from polyacrylamide gel electrophoresis as to material length.

EXAMPLE 74

10 Synthesis of Sequence Specific Pyrrolidine Oligomer Having Phosphodiester Linkages

"Aforvirsen" is an anti-papilloma agent having the nucleobase sequence:

TTG CTT CCA TCT TCC TCG TC.

A pyrrolidine phosphodiester linked oligomer of this preselected sequence is prepared using the T, A, C and G reagents from Examples 7, 11, 13 and 14, respectively, as per the procedure of Example 73 using iodine as the oxidation reagent to give the phosphodiester linked oligomeric compound having the "Aforvirsen" sequence.

20

EXAMPLE 75

Synthesis of Sequence Specific Pyrrolidine Oligomer Having Phosphorothioate Linkages

A pyrrolidine phosphorothioate-linked oligomer of sequence TTG CTT CCA TCT TCC TCG TC is prepared using the T, A, C and G reagents from Examples 7, 11, 13 and 14, respectively, as per the procedure of Example 73 using 3H-1,2-benzodithiole-3-one 1,1-dioxide as the oxidation reagent to give the phosphorothioate linked oligomeric compound.

30 Example 76

Preparation of N-(2-[1-Tyrosinyl]-acetyl)-2-hydroxymethyl-pyrrolidine-4-morpholinophosphoramidate-thymidine-

- 90 -

phosphodiester-thymidine Trimer.

A controlled pore glass resin derivatized with a 3'-5' phosphodiester linked dimer of thymidine [5'-DMT-O-T-O-P(=O)-O-T-CPG] was synthesized in a standard manner. Two aliquots of this resin (each 24.6 mg, 47.5 mmole/g resin) were separately detritylated using 3 % trichloroacetic acid in dichloromethane and then sequentially washed with anhydrous acetonitrile and pyridine. Each was vigorously agitated with 1 mL of a solution of the N-(2-[1-Tyrosinyl]acetyl)-2-O-dimethoxytrityloxymethyl-pyrrolidine-3-H-phosphonate monomer (25 mM) in adamantoyl chloride (50 mM) and pyridine for five minutes. The resins were then washed repeatedly with anhydrous pyridine and blown dry with argon gas. To one column was added 1 mL of iodine/pyridine/water (2/90/8) solution and agitated vigorously; to the second column was added 1 mL of morpholine/carbon tetrachloride/pyridine (1/5/5) solution and both were agitated for 30 minutes. The columns were subsequently washed with pyridine and acetonitrile and then blown dry with argon gas. Products were cleaved from the CPG by treatment with 1 mL of concentrated ammonium hydroxide for 30 minutes. The ammonia solutions were evaporated to dryness and then treated with 80 % acetic acid in water for 30 minutes to remove the trityl group. An aliquot of the solution from the first column exhibited a single peak in an HPLC analysis (27.8 min, Vyadec C-18, 260 nm, linear gradient of ammonium acetate (pH 7)-acetonitrile, 0-75 % acetonitrile in 52 minutes) corresponding to the phosphodiester trimer. A similar analysis yielded a pair of HPLC peaks (39.7 and 39.9 min) corresponding to the phosphoramidate diastereomeric trimers.

In a parallel set of experiments using the HPP-Tyrosine H-phosphonate monomer and the 5'-DMT-O-T-O-P(=O)-O-T-CPG, the coupled resins were oxidized to the phosphodiester trimer or to the phosphoramidate trimer using dimethylamine/carbon tetrachloride. These experiments yielded the phosphodiester

trimer and the unresolved dimethylamine phosphoramidate trimers at 12.4 min by an HPLC analysis, under the conditions described before.

EXAMPLE 77**5 (5R)-O-(t-Butyldimethylsilyl)-(3R)-Hydroxypiperidine**

N-Benzyl-(5R)-O-(t-butyldimethylsilyl)-(3R)-hydroxypiperidine (J. Cossy, C. Dumas, P. Michel, D. Gomez Pardo, *Tetrahedron Lett.*, 1995, 36, 549) is dissolved in ethanol. A catalytic amount of 10% Pd/C is added, and the solution
10 shaken under 3 ATM H₂. Once the starting material is consumed, the catalyst is filtered and the solvent removed under reduced pressure. The product is used without further purification.

EXAMPLE 78**15 N-(Fmoc)-(5R)-O-(t-Butyldimethylsilyl)-(3R)-Hydroxypiperidine**

(5R)-O-(t-butyldimethylsilyl)-(3R)-hydroxypiperidine and Na₂CO₃ (1.3 eq) are suspended in H₂O/Dioxane (1:1) (0.1M). Fluorenylmethyl chloroformate (1.1 eq) in toluene is added dropwise. The temperature of the reaction is not allowed to
20 rise above 25 °C during the addition. The mixture is stirred vigorously overnight, acidified to pH 3 with concentrated HCl, and extracted twice with ethyl acetate. The organic extract is washed with brine. The solution is dried with MgSO₄, filtered and the solvent removed in vacuo. The product
25 is purified by silica gel flash column chromatography.

EXAMPLE 79**N-(Fmoc)-(3R)-O-Dimethoxytrityl-(5R)-O-(t-Butyldimethylsilyl)-Piperidine**

N-(Fmoc)-(5R)-O-(t-butyldimethylsilyl)-(3R)-hydroxypiperidine is coevaporated with dry pyridine, and
30 redissolved in dry pyridine (0.1M). Dimethoxytrityl chloride

- 92 -

(1.2 eq) is added in portions over 15 minutes, and the solution stirred at RT overnight. Methanol is then added (10 ml), and the solvent removed under reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, NaHCO₃, brine, dried with MgSO₄, and evaporated. The product is purified by silica gel flash column chromatography.

EXAMPLE 80**N-(Fmoc)-(3R)-O-Dimethoxytrityl-(5R)-Hydroxypiperidine**

N-(Fmoc)-(3R)-O-dimethoxytrityl-(5R)-O-(t-butyldimethylsilyl)-Piperidine is dissolved in THF (0.1M), and added to a solution of tetrabutylammonium fluoride (3 eq) and acetic acid (9 eq) in THF. The solution is stirred until the starting material is consumed. The reaction is quenched with NaHCO₃, extracted with diethyl ether, washed with NaHCO₃, brine and dried with MgSO₄.

EXAMPLE 81**N-(Fmoc)-(3R)-O-Dimethoxytrityl-Piperidine-(5R)-O-Hydrogen Phosphonate Triethylammonium salt**

The title compound is prepared from N-(Fmoc)-(3R)-O-dimethoxytrityl-(5R)-hydroxypiperidine using the general procedure of Example 59.

EXAMPLE 82**N-(Fmoc)-(3R)-O-Dimethoxytrityl-Piperidine-(5R)-O-Succinate**

The title compound is prepared from N-(Fmoc)-(3R)-O-dimethoxytrityl-(5R)-hydroxypiperidine using the general procedure of Example 60.

EXAMPLE 83**N-(Fmoc)-(3R)-O-Dimethoxytrityl-Piperidine-(5R)-O-Succinyl-**

30 CPG

The title compound is prepared from N-(Fmoc)-(3R)-O-dimethoxytrityl-piperidine-(5R)-O-succinate using the general procedure of Example 61.

EXAMPLE 84

5 **(2S)-Carboxyethyl-Pyrrolidine-5-carboxylic acid-4-methylphenylthioester**

The title compound is obtained as a mixture of epimers at C5 from N-BOC-Pyroglutamate ethyl ester using the method of J. Ezquerra, A., Rubio, C., Pedregal, G., Sanz, J. H.,
10 Rodriguez, J. L., Garcia R., *Tetrahedron Lett.* **1993**, *34*, 4989.

EXAMPLE 85

N-Fmoc-(2S)-Carboxyethyl-Pyrrolidine-5-carboxylic acid-4-methylphenylthioester

15 (2S)-Carboxyethyl-pyrrolidine-5-carboxylic acid-4-methylphenylthioester and Na₂CO₃ (1.3 eq) are suspended in H₂O/Dioxane (1:1) (0.1 M). Fluorenylmethyl chloroformate (1.1 eq) in toluene is added dropwise. The temperature of the reaction is not allowed to rise above 25 °C during the
20 addition. The mixture is stirred vigorously overnight, acidified to pH 3 with concentrated HCl, and extracted twice with ethyl acetate. The organic extract is washed with brine. The solution is dried with MgSO₄, filtered and the solvent removed *in vacuo*. The product is purified by silica
25 gel flash column chromatography.

EXAMPLE 86

**N-Fmoc-(2S)-Carboxyethyl-(5R)-Hydroxymethyl-Pyrrolidine and
N-Fmoc-(2S)-Carboxyethyl-(5S)-Hydroxymethyl-Pyrrolidine**

The mixture of thioesters from the Example 85 are
30 dissolved in THF and treated with Pd(OAc)₂ and triethylsilane. Once the thioesters are reduced to the aldehyde, the solution

- 94 -

is cooled to 0 °C, and 1 eq LiBH₄ added. The reaction is stopped by the addition of acetic acid. The solvent is removed, and the residue extracted with ethyl acetate, washed with 5% NaHCO₃, brine and dried with MgSO₄. The products
5 (epimers) are separated by silica gel flash column chromatography.

EXAMPLE 87**N-FMOC-(2S)-Carboxyethyl-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine**

10 N-FMOC-(2S)-carboxyethyl-(5R)-hydroxymethyl-pyrrolidine is coevaporated with dry pyridine, and redissolved in dry pyridine (0.1M). Dimethoxytrityl chloride (1.2 eq) is added in portions over 15 minutes, and the solution stirred at RT overnight. Methanol is then added (10 ml), and the solvent
15 removed under reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, NaHCO₃, brine, dried with MgSO₄, and evaporated. The residue is purified by silica gel flash column chromatography.

EXAMPLE 88

20 **N-FMOC-(2S)-Hydroxymethyl-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine**

N-FMOC-(2S)-carboxyethyl-(5R)-dimethoxytrityloxymethyl-pyrrolidine is dissolved in dry THF, cooled to 0 °C, and 2 eq LiBH₄ is added. The solution is warmed to room temperature
25 until the starting material is consumed. The reaction is quenched with ethyl acetate, and washed with 0.1 M citric acid, NaHCO₃, brine, dried with MgSO₄ and evaporated. The product is purified by silica gel flash column chromatography.

30 **EXAMPLE 89**

N-FMOC-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-

- 95 -

Oxymethyl-Hydrogen Phosphonate Triethylammonium salt

The title product is prepared from N-FMOC-(2S)-Hydroxymethyl-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine using the general procedure of Example 59.

5 **EXAMPLE 90****N-FMOC-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-Oxymethyl Succinate**

The title product is prepared from N-FMOC-(2S)-Hydroxymethyl-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine using
10 the general procedure of Example 60.

EXAMPLE 91**N-FMOC-(5R)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-Oxymethyl-Succinyl-CPG**

The title product is prepared from N-FMOC-(5R)-
15 dimethoxytrityloxymethyl-pyrrolidine-(2S)-Oxymethyl Succinate using the general procedure of Example 61.

EXAMPLE 92**N-FMOC-(2S)-Carboxyethyl-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine**

20 N-FMOC-(2S)-Carboxyethyl-(5S)-Hydroxymethyl-Pyrrolidine is coevaporated with dry pyridine, and redissolved in dry pyridine (0.1M). Dimethoxytrityl chloride (1.2 eq) is added in portions over 15 minutes and the solution is stirred at RT overnight. Methanol (10 ml) is then added and the solvent
25 removed under reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, concentrated NaHCO₃ solution, brine, dried with MgSO₄, and evaporated. The residue is purified by silica gel flash column chromatography.

30 **EXAMPLE 93**

- 96 -

N-FMOC-(2S)-Hydroxymethyl-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine

N-FMOC-(2S)-carboxyethyl-(5S)-dimethoxytrityloxymethyl-pyrrolidine is dissolved in dry THF, cooled to 0 °C, and 2 eq
5 LiBH₄ is added. The solution is warmed to room temperature until the starting material is consumed. The reaction is quenched with ethyl acetate, and washed with 0.1 M citric acid, concentrated NaHCO₃ solution, brine, dried with MgSO₄ and evaporated. The product is purified by silica gel flash
10 column chromatography.

EXAMPLE 94

N-FMOC-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-oxymethyl-Hydrogen Phosphonate Triethylammonium salt

The title product is prepared from N-FMOC-(2S)-
15 hydroxymethyl-(5S)-dimethoxytrityloxymethyl-pyrrolidine using the general procedure of Example 59.

EXAMPLE 95

N-FMOC-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-Oxymethyl Succinate

20 The title product is prepared from N-FMOC-(2S)-Hydroxymethyl-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine fusing the general procedure of Example 60.

EXAMPLE 96

N-FMOC-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-Oxymethyl-Succinyl-CPG

25 The title product is prepared from N-FMOC-(5S)-Dimethoxytrityloxymethyl-Pyrrolidine-(2S)-Oxymethyl Succinate using the general procedure of Example 61.

EXAMPLE 97

N-FMOC-2,2'-Dihydroxyethylamine

30

- 97 -

Diethanolamine and Na_2CO_3 (1.3 eq) are suspended in H_2O /Dioxane (1:1) (0.1M). Fluorenylmethyl chloroformate (1.1 eq) in toluene is added dropwise. The temperature of the reaction is not allowed to rise above 25 °C during the addition. The mixture is stirred vigorously overnight, acidified to pH 3 with concentrated HCl, and extracted twice with ethyl acetate. The organic extract is washed with brine, dried with MgSO_4 , filtered and the solvent removed in vacuo. The product is purified by silica gel flash column chromatography.

EXAMPLE 98**N-FMOC-2'-Hydroxyethyl-2-O-Dimethoxytritylethylamine**

N-FMOC-2,2'-dihydroxyethylamine is coevaporated with dry pyridine, and redissolved in dry pyridine (0.1 M). Dimethoxytrityl chloride (1.2 eq) is added in portions over 15 minutes, and the solution stirred at RT overnight. Methanol (10 ml) is then added and the solvent removed under reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, concentrated NaHCO_3 solution, brine, dried with MgSO_4 , and evaporated. The residue is purified by silica gel flash column chromatography.

EXAMPLE 99**N-FMOC-2-O-Dimethoxytritylethylamino-2'-O-Ethyl Hydrogen Phosphonate Triethylammonium salt**

The title product is prepared from N-FMOC-2'-Hydroxyethyl-2-O-Dimethoxytritylethylamine using the general procedure of Example 59.

EXAMPLE 100**N-FMOC-2,3-Propanediol**

1-Aminopropanediol and Na_2CO_3 (1.3 eq) are suspended in

- 98 -

H₂O/Dioxane (1:1) (0.1 M). Fluorenylmethyl chloroformate (1.1 eq) in toluene is added dropwise. The temperature of the reaction is not allowed to rise above 25 °C during the addition. The mixture is stirred vigorously overnight, acidified to pH 3 with concentrated HCl, and extracted twice with ethyl acetate. The organic extract is washed with brine, dried with MgSO₄, filtered and the solvent removed in vacuo. The product is purified by silica gel flash column chromatography.

10 EXAMPLE 101

N-FMOC-2-Hydroxy-3-O-Dimethoxytrityl-Propane

N-FMOC-2,3-Propanediol is coevaporated with dry pyridine, and redissolved in dry pyridine (0.1 M). Dimethoxytrityl chloride (1.2 eq) is added in portions over 15 minutes and the solution stirred at RT overnight. Methanol (10 ml) is then added and the solvent removed under reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, concentrated NaHCO₃ solution, brine, dried with MgSO₄, and evaporated. The 20 residue is purified by silica gel flash column chromatography.

EXAMPLE 102

Derivatization of Backbone Segment Amino Combinatorial Site to form Ureas General Procedure

25 To the free amine on solid support is added a solution of carbonyl diimidazole (0.1 mmol/ml) in DMF. A tenfold excess of reagent is added at the 10 umol scale level. The reaction is allowed to proceed for 30 min, and the reagent removed with DMF. A solution of the desired amine in DMF 30 (10% v/v) is then added. After 30 minutes the solid support is washed with DMF until all the reagents are removed to give the urea derivatized backbone segment amino combinatorial

- 99 -

site.

EXAMPLE 103**N-Fmoc-3,4-Trans-Dicarboxymethylpyrrolidine**

Dimethylfumarate (38.7 g, 268 mmol) was dissolved in 1 l
5 CH₂Cl₂, and trifluoroacetic acid added (3 ml, 40 mmol). The
Azomethine ylide precursor N-Benzyl-N-Methoxymethyl-N-
(Trimethylsilylmethyl)-amine was added dropwise with stirring
(60.4 g, 255 mmol). The solution was stirred overnight at
RT. The solvent was removed and the crude product was
10 redissolved in 1.5 l methanol. 300 ml 1 N HCl was added, the
solution degassed with Ar, and 10 g 10% Pd/C added. H₂ was
bubbled through then the solution stirred under a H₂ balloon
until the material was consumed (24 hr). The catalyst was
filtered, the solvent removed and the residue redissolved in
15 500 ml H₂O and 500 ml dioxane to which was added 32 g (300
mmol) Na₂CO₃. A solution of Fmoc-Cl (66 g, 255 mmol) in 500
ml dioxane was added dropwise. The solution was stirred 16
h, acidified to pH 3 and the solvent evaporated. The residue
was extracted with EtOAc, washed with water, brine dried
20 with MgSO₄ and evaporated. 92 g title product was obtained.

EXAMPLE 104**N-Fmoc-3,4-Trans-Dihydroxymethylpyrrolidine**

N-Fmoc-3,4-Trans-Dicarboxymethylpyrrolidine (92 g, 225 mmol)
was dissolved in 1000 ml THF, and 250 ml LiBH₄ (2M in THF,
25 Aldrich) was added with cooling (exothermic). The solution
was stirred at RT 2 hrs and carefully quenched with 50% HCl
to pH 2. The solvent was evaporated. The residue was
extracted with EtOAc (2 X 500 ml), washed with NaHCO₃, brine,
and dried with MgSO₄ and evaporated. 78.6 g crude product was
30 obtained (100 %). The product was recrystallized from 300 ml
acetonitrile to give 60 g

- 100 -

EXAMPLE 105**N-Fmoc-3-Dimethoxytrityloxymethyl-4-hydroxymethylpyrrolidine**

N-Fmoc-3,4-Trans-Dihydroxymethylpyrrolidine was dissolved in CH_2Cl_2 and pyridine (9:1), cooled to 0 °C and DMT-Cl added in portions (1.0 eq). The solution was stirred overnight at 0 °C. The reaction was quenched with MeOH, evaporated and excess solvent removed by azeotropic distillation with toluene. The product was redissolved in EtOAc, washed with 5% citric acid, NaHCO_3 , brine and dried with MgSO_4 .

Purification was achieved by silica gel flash column chromatography, using a gradient of from 0 to 3% methanol, in CH_2Cl_2 . The product was obtained in 50% yield, along with 25% ditritylated material which is conserved and recycled, and 25% unreacted starting material.

EXAMPLE 106**N-FMOC-trans-3-(oxymethyl)-4-(Dimethoxytrityloxymethyl)-Pyrrolidine Hydrogen Phosphonate Triethylammonium salt**

The title product is prepared from N-Fmoc-3-Dimethoxytrityloxymethyl-4-hydroxymethylpyrrolidine using the general procedure of Example 59.

EXAMPLE 107**N-FMOC-trans-3-(oxymethyl)-4-(Dimethoxytrityloxymethyl)-Pyrrolidine Succinate**

The title product is prepared from N-FMOC-trans-3-(Hydroxymethyl)-4-(Dimethoxytrityloxymethyl)-Pyrrolidine using the general procedure of Example 60.

EXAMPLE 108**N-FMOC-trans-3-(oxymethyl)-4-(Dimethoxytrityloxymethyl)-Pyrrolidine Succinyl CPG**

The title product is prepared from N-FMOC-trans-3-(Hydroxymethyl)-4-(Dimethoxytrityloxymethyl)-Pyrrolidine

- 101 -

using the general procedure of Example 61.

EXAMPLE 109**N-CBZ-cis-4-Amino-2-cyclopenten-1-ol**

The product is prepared by the method of A. R. Ritter,
5 M. J. Miller, Tetrahedron Lett. 1994, 35, 9379.

EXAMPLE 110**N-CBZ-cis-4-Amino-2,3-oxocyclopentan-1-ol**

N-CBZ-cis-4-Amino-2-cyclopenten-1-ol is dissolved in
dichloromethane and meta chloroperbenzoic acid is added. The
10 solution is stirred until the starting material disappears.
The reaction is quenched by the addition of a 5% solution of
sodium bisulfite, extracted with dichloromethane, washed with
NaHCO₃, brine and dried. The title compound is purified by
silica gel flash column chromatography.

EXAMPLE 111**N-CBZ-cis-4-Aminocyclopentan-1,2-diol**

The epoxide of Example 110 is dissolved in THF, and
treated with 2 eq LiBH₄ in the presence of a catalytic amount
of Ti(OiPr)₄. Stirring is continued until the reaction is
20 complete. The reaction is diluted with ethyl acetate, washed
with 0.1 M citric acid, NaHCO₃, brine, dried with MgSO₄ and
evaporated. The title compound is purified by silica gel
flash column chromatography.

EXAMPLE 112**N-CBZ-cis-4-Amino-2-O-dimethoxytritylcyclopentan-1-ol**

N-CBZ-cis-4-Aminocyclopentan-1,2-diol is coevaporated
with dry pyridine, and redissolved in dry pyridine (0.1 M).
Dimethoxytrityl chloride (1.0 eq) is added in portions over
15 minutes and the solution stirred at RT overnight.
30 Methanol is then added (10 ml), and the solvent removed under

- 102 -

reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, NaHCO_3 , brine, dried with MgSO_4 , and evaporated. The title compound is purified by silica gel flash column chromatography.

5 **EXAMPLE 113****N-FMOC-cis-4-Amino-2-O-dimethoxytritylcyclopentan-1-ol**

N-CBZ-cis-4-Amino-2-O-dimethoxytritylcyclopentan-1-ol is dissolved in ethanol and 10% Pd/C is added. The mixture is shaken under 1 ATM H_2 until all the material is consumed. The catalyst is filtered, the solvent removed. The residue and Na_2CO_3 (1.3 eq) are suspended in H_2O /Dioxane (1:1) (0.1 M). Fluorenylmethyl chloroformate (1.1 eq) in toluene is added dropwise. The temperature of the reaction is not allowed to rise above 25 °C during the addition. The mixture is stirred vigorously overnight, acidified to pH 3 with concentrated HCl, and extracted twice with ethyl acetate. The organic extract is washed with brine, dried with MgSO_4 , filtered and the solvent removed in vacuo. The title compound is purified by silica gel flash column chromatography.

20 **EXAMPLE 114****N-FMOC-cis-4-Amino-2-O-dimethoxytritylcyclopentan-1-O-Hydrogen Phosphonate triethylammonium salt**

The title compound is prepared using the general procedure of Example 59.

25 **EXAMPLE 115****N-CBZ-cis-4-Amino-2-cyclohexen-1-ol**

The product is prepared by the method of A. R. Ritter, M. J. Miller, J. Org. Chem. 1994, 59, 4602.

EXAMPLE 11630 **N-CBZ-cis-4-Amino-2,3-oxocyclohexan-1-ol**

- 103 -

N-CBZ-cis-4-Amino-2-cyclohexen-1-ol is dissolved in dichloromethane and meta chloroperbenzoic acid is added. The solution is stirred until the starting material disappears. The reaction is quenched by the addition of a 5% solution of sodium bisulfite, extracted with dichloromethane, washed with NaHCO₃, brine and dried. The title compound is purified by silica gel flash column chromatography.

EXAMPLE 117**N-CBZ-cis-4-Aminocyclohexan-1,2-diol**

The epoxide of the previous example is dissolved in THF, and treated with 2 eq LiBH₄ in the presence of a catalytic amount of Ti(OiPr)₄. Stirring is continued until the reaction is complete. The reaction is diluted with ethyl acetate, washed with 0.1M citric acid, NaHCO₃, brine, dried with MgSO₄, and evaporated. The title compound is purified by silica gel flash column chromatography.

EXAMPLE 118**N-CBZ-cis-4-Amino-2-O-dimethoxytritylcyclohexan-1-ol**

N-CBZ-cis-4-aminocyclohexan-1,2-diol is coevaporated with dry pyridine, and redissolved in dry pyridine (0.1 M). Dimethoxytrityl chloride (1.0 eq) is added in portions over 15 min, and the solution stirred at RT overnight. Methanol is then added (10 ml), and the solvent removed under reduced pressure. The resulting gum is redissolved in ethyl acetate, washed with 0.1 M citric acid, NaHCO₃, brine, dried with MgSO₄, and evaporated. The title compound is purified by silica gel flash column chromatography.

EXAMPLE 119**N-FMOC-cis-4-Amino-2-O-dimethoxytritylcyclohexan-1-ol**

N-CBZ-cis-4-Amino-2-O-dimethoxytritylcyclohexan-1-ol is dissolved in ethanol and 10% Pd/C is added. The mixture is

- 104 -

shaken under 1 ATM H₂ until all the material is consumed. The catalyst is filtered, the solvent removed. The residue and Na₂CO₃ (1.3 eq) are suspended in H₂O/Dioxane (1:1) (0.1 M). Fluorenylmethyl chloroformate (1.1 eq) in toluene is added dropwise. The temperature of the reaction is not allowed to rise above 25 °C during the addition. The mixture is stirred vigorously overnight, acidified to pH 3 with concentrated HCl, and extracted twice with ethyl acetate. The organic extract is washed with brine, dried with MgSO₄, filtered and the solvent removed in vacuo. The title compound is purified by silica gel flash column chromatography.

EXAMPLE 120**N-Fmoc-cis-4-Amino-2-O-dimethoxytritylcyclohexan-1-O-Hydrogen Phosphonate triethylammonium salt**

The title compound is prepared using the general procedure of Example 59.

Example 121

Synthesis of a 4-mer having backbone segment amino combinatorial Backbone Segment Amino Combinatorial Sites combinatorialized with the aldehydes benzaldehyde, Aldrich-B133-4; m-tolualdehyde, Aldrich-T3,550-5; m-anisaldehyde, Aldrich-12,965-8; and 3-nitrobenzaldehyde, Aldrich-N1,084-5

Method 1: Bead Splitting

A solid support is derivatized with 2-O-(dimethoxytrityl)ethylsuccinate half ester as in Example 26. The DMT protecting group is removed using the standard method of Example 64. 4-O-Dimethoxytrityl-2-N-Fmoc-aminobutan-1-ol is treated with PCl₃ as per Example 28 to form the phosphonic acid which is condensed onto the derivatized resin as in Example 38. The Fmoc amino protecting group is removed as per Example 4 and the solid support is divided into 4 equal portions and each portion is reacted with one of benzalde-

- 105 -

hyde, m-tolualdehyde, m-anisaldehyde, or 3-nitrobenzaldehyde to effect coupling to the resulting free amino as per the method of Example 66 and the method of Look, G. C., et al., *Tetrahedron Lett.*, 1995, 36, 2937-2940, and the portions of solid support recombined. The combined solid support is treated with a solution of NaCNBH_3 or LiBH_4 in tetrahydrofuran for 30 minutes followed by washing of the solid support by methanol. The above methods are repeated until four aminodiol monomer subunits are incorporated having an equal molar mixture of each aldehyde letter at each of the amino sites. The resulting four mer is treated using the procedures of Example 67, with a large excess of benzylamine in carbon tetrachloride/pyridine. The solid support is shaken for 15 minutes and the supernatant is removed by filtration and then washed with pyridine. A second treatment with a large excess of benzylamine in carbon tetrachloride/pyridine followed by shaking will insure efficient oxidation to the phosphoramidate. The resin is washed with pyridine/acetonitrile and then the phosphoramidate is cleaved from the resin by a treatment with concentrated ammonium hydroxide at room temperature for 3 hours. Evaporation of the supernatant and purification of the phosphoramidate on an RP-18 HPLC column will yield the final oligomer.

Method 2

A solid support is derivatized with 2-O-(dimethoxytrityl)ethylsuccinate half ester as in Example 26. The DMT protecting group is removed using the standard method of Example 64. 4-O-Dimethoxytrityl-2-N-Fmoc-aminobutan-1-ol is treated with PCl_3 as per Example 28 to form the phosphonic acid which is condensed onto the derivatized resin as in Example 38. Following this procedure the four mer is synthesized in four iterations of the above. The Fmoc amino protecting groups are removed as per Example 4 and is reacted

- 106 -

concurrently, in one pot, with benzaldehyde, m-tolualdehyde, m-anisaldehyde, and 3-nitrobenzaldehyde. To effect this concurrent reaction 50 μ mol of oligomeric compound attached to solid support is reacted with 50 μ mol benzaldehyde, 50 μ mol m-tolualdehyde, 50 μ mol m-anisaldehyde, and 50 μ mol 3-nitrobenzaldehyde in triethyl orthoformate. The resulting imines are reduced as per the procedure of method 1 above. In this way the four amino combinatorial sites of each oligomeric compound attached to the solid support are combinatorialized utilizing a competitive mechanism involving a mixture of aldehydes. The resulting four mer is treated using the procedures of Example 67, with a large excess of benzylamine in carbon tetrachloride/pyridine. The solid support is shaken for 15 minutes and the supernatant is removed by filtration and then washed with pyridine. A second treatment with a large excess of benzylamine in carbon tetrachloride/ pyridine followed by shaking will insure efficient oxidation to the phosphoramidate. The resin is washed with pyridine/acetonitrile and then the phosphoramidate is cleaved from the resin by a treatment with concentrated ammonium hydroxide at room temperature for 3 hours. Evaporation of the supernatant and purification of the phosphoramidate on an RP-18 HPLC column will yield the final oligomer.

EXAMPLE 122

Synthesis of Combinatorial libraries using various selected aldehydes

Using the procedure of example 121 libraries are prepared from oligomeric compounds of the invention that are derivatized with one, two, three, four or more of the following aldehydes available from Aldrich Chemical Company, Inc., Milwaukee, WI. The Aldrich catalog number is given in the right hand column and the compound name is given in the

- 107 -

left hand column:

Aromatic aldehydes

	10793-5	Phenylacetaldehyde
	D20425	Diphenylacetaldehyde
5	24582-8	Hydrocinnamaldehyde
	24136-9	Phenylpropionaldehyde
	28902-7	(+/-)-3-Phenylbutyraldehyde
	28899-3	Alpha-amylcinnamaldehyde
	16116-0	Alpha-bromocinnamaldehyde
10	26813-5	4-Stilbenecarboxaldehyde
	B133-4	Benzaldehyde
	11755-2	o-Tolualdehyde
	25069-4	Alpha.alpha.alpha-trifluoro-o-tolualdehyde
	F480-7	2-Fluorobenzaldehyde
15	12497-4	2-Chlorobenzaldehyde
	B5700-1	2-Bromobenzaldehyde
	10962-2	o-Anisaldehyde
	15372-9	2-Ethoxybenzaldehyde
	N1080-2	2-Nitrobenzaldehyde
20	T3550-5	m-Tolualdehyde
	19687-8	Alpha.alpha.alpha-trifluoro-m-tolualdehyde
	F500-5	3-Fluorobenzaldehyde
	C2340-3	3-Chlorobenzaldehyde
	B5720-6	3-Chlorobenzaldehyde
25	12965-8	m-Anisaldehyde
	34648-9	3-(Trifluoromethoxy)-benzaldehyde
	34199-1	3-(1,1,2,2-Tetrafluoroethoxy)-benzaldehyde
	H1980-8	3-Hydroxybenzaldehyde
	N1084-5	3-Nitrobenzaldehyde
30	11528-2	Isophthalaldehyde
	T3560-2	p-Tolualdehyde
	23363-3	4-Ethylbenzaldehyde
	13517-8	4-Isopropylbenzaldehyde
	22494-4	Alpha.alpha.alpha-trifluoro-p-tolualdehyde
35	12837-6	4-Fluorobenzaldehyde

	11221-6	4-Chlorobenzaldehyde
	B5740-0	4-Bromobenzaldehyde
	A8810-7	p-Anisaldehyde
	17360-6	4-Ethoxybenzaldehyde
5	33363-8	4-Propoxybenzaldehyde
	23808-2	4-Butoxybenzaldehyde
	37060-6	4-(Trifluoromethoxy)-benzaldehyde
	27486-0	Terephthaldehyde mono-(diethyl acetal)
	14408-8	4-Hydroxybenzaldehyde
10	22277-1	4-(Methylthio)benzaldehyde
	10976-2	4-(Dimethylamino)benzaldehyde
	D8625-6	4-(Dimethylamino)benzaldehyde
	33851-6	4-(Dibutylamino)benzaldehyde
	29355-5	4-(3-Dimethylaminopropoxy)benzaldehyde
15	13017-6	4-Nitrobenzaldehyde
	T220-7	Terephthalaldicarboxaldehyde
	34252-1	3-Fluoro-2-methylbenzaldehyde
	34649-7	2-Fluoro-3-(trifluoromethyl)-benzaldehyde
	26514-4	2,3-Difluorobenzaldehyde
20	26515-2	2,6-Difluorobenzaldehyde
	14124-0	2-Chloro-6-fluorobenzaldehyde
	D5650-0	2,6-Dichlorobenzaldehyde
	25483-5	2,3-Dichlorobenzaldehyde
	D13020-6	2,3-Dimethoxybenzaldehyde
25	29250-8	2,6-Dimethoxybenzaldehyde
	31980-5	3-Fluorosalicylaldehyde
	12080-4	o-Vanillin
	18983-9	2,3-Dihydroxybenzaldehyde
	10604-6	2-Chloro-6-nitrobenzaldehyde
30	16382-1	3-methoxy-2-nitrobenzaldehyde
	11750-1	2,6-Dinitrobenzaldehyde
	15104-1	2,4-Dimethylbenzaldehyde
	15106-8	2,5-Dimethylbenzaldehyde
	37682-5	2-Chloro-5-(trifluoromethyl)benzaldehyde
35	26516-0	3,4-Difluorobenzaldehyde

	26517-9	2,4-Difluorobenzaldehyde
	26518-7	2,5-Difluorobenzaldehyde
	30600-2	3-Chloro-4-fluorobenzaldehyde
	34807-4	2-Chloro-4-fluorobenzaldehyde
5	33954-7	3-Bromo-3-fluorobenzaldehyde
	D5660-8	3,4-Dichlorobenzaldehyde
	14675-7	2,4-Dichlorobenzaldehyde
	15212-9	3-Methyl- <i>p</i> -anisaldehyde
	15558-6	3-Fluoro- <i>p</i> -anisaldehyde
10	15429-6	5-Bromo- <i>o</i> -anisaldehyde
	D13040-0	2,4-Dimethoxybenzaldehyde
	D13060-5	2,5-Dimethoxybenzaldehyde
	14375-8	3,4-Dimethoxybenzaldehyde
	25275-1	3-Ethoxy-4-methoxybenzaldehyde
15	P4910-4	Piperonal
	26459-8	1,4-Benzodioxan-6-carboxaldehyde
	31691-1	4-Hydroxy-3-methylbenzaldehyde
	34606-3	2-Chloro-4-hydroxybenzaldehyde
	25975-6	5-Chlorosalicylaldehyde
20	13728-6	5-Bromosalicylaldehyde
	14686-2	2-Hydroxy-5-methoxybenzaldehyde
	16069-5	2-Hydroxy-4-methoxybenzaldehyde
	14368-5	3-Hydroxy-4-methoxybenzaldehyde
	V110-4	Vanillin
25	12809-0	3-Ethoxy-4-hydroxybenzaldehyde
	34215-7	5-(Trifluoromethoxy) salicylaldehyde
	D10840-5	3,4-Dihydroxybenzaldehyde
	D10820-0	2,5-Dihydroxybenzaldehyde
	16863-7	2,4-Dihydroxybenzaldehyde
30	22568-1	4-(Diethylamino) salicylaldehyde
	C5880-0	5-Chloro-2-nitrobenzaldehyde
	13903-3	2-Chloro-5-nitrobenzaldehyde
	C5870-3	4-Chloro-3-nitrobenzaldehyde
	14432-0	4-Hydroxy-3-nitrobenzaldehyde
35	15616-7	3-Hydroxy-4-nitrobenzaldehyde

- 110 -

	27535-2	2-Hydroxy-5-nitrobenzaldehyde
	H4810-7	5-Hydroxy-2-nitrobenzaldehyde
	D19360-7	2,4-Nitrobenzaldehyde
	29013-0	3,5-Bis(trifluoromethyl)benzaldehyde
5	29017-3	3,5-Difluorobenzaldehyde
	13940-8	3,5-Dichlorobenzaldehyde
	36811-3	3,5-Dihydroxybenzaldehyde
	12269-2	3,5-Dimethoxybenzaldehyde
	36810-5	3,5-Dibenzyloxybenzaldehyde
10	M680-8	Mesitaldehyde
	29233-8	2,3,5-Trichlorobenzaldehyde
	13061-3	5-Bromoveratraldehyde
	13871-1	2,4,6-Trimethoxybenzaldehyde
	T6840-3	3,4,5-Trimethoxybenzaldehyde
15	14039-2	3,5-Dimethyl-4-hydroxybenzaldehyde
	35768-5	2,6-Dimethyl-4-hydroxybenzaldehyde
	14040-6	3,5-Di-tert-butyl-4-hydroxybenzaldehyde hemihydrate
	26181-5	3,5-Dichlorosalicylaldehyde
20	12213-0	3,5-Dibromosalicylaldehyde
	28344-4	3,5-Diiodosalicylaldehyde
	13060-5	5-Bromovanillin
	12948-8	5-Iodovanillin
	13879-7	4,6-Dimethoxysalicylaldehyde
25	25871-7	5-Nitrovanillin
	S760-2	3,5-Dinitrosalicylaldehyde
	25959-4	2,5-Dimethyl-p-anisaldehyde
	T6540-4	5-Bromo-2,4-dimethoxybenzaldehyde
	N2800-0	4-Nitrovanillin
30	27680-4	3,5-Dinitrosalicylaldehyde
	15205-6	2,5-Dimethyl-p-anisaldehyde
	29251-6	5-Bromo-2,4-dimethoxybenzaldehyde
	15557-8	6-Bromoveratraldehyde
	13215-2	2,4,5-Trimethoxybenzaldehyde
35	27960-9	6-Nitroveratraldehyde

- 111 -

	13765-0	6-Nitropiperonal
	27679-0	2,5-Dichloroterephthaldehyde
	33066-3	2,3,4-Trifluorobenzaldehyde
	29231-1	2,3,6-Trichlorobenzaldehyde
5	15201-3	2,3-Dimethyl- <i>p</i> -anisaldehyde
	29627-9	2,4-Dimethoxy-3-methylbenzaldehyde
	15209-9	2,3,4-Trimethoxybenzaldehyde
	26084-3	2,3,4-Trihydroxybenzaldehyde
	32893-6	Tetrafluorobenzaldehyde
10	10374-8	Pentafluorobenzaldehyde
	B3468-0	4-Biphenylcarboxaldehyde
	19175-2	3-Phenoxybenzaldehyde
	B2700-5	3-Benzloxybenzaldehyde
	19540-5	3-(4-Methylphenoxy)benzaldehyde
15	19592-8	3-(4- <i>tert</i> -Butylphenoxy)benzaldehyde
	19539-1	3-[3-(Trifluoromethyl)phenoxy]benzaldehyde
	19530-8	3-(4-Chlorophenoxy)benzaldehyde
	19590-1	3-(3,4-Dichlorophenoxy)benzaldehyde
	19774-2	3-(3,5-Dichlorophenoxy)benzaldehyde
20	19589-8	3-(4-Methoxyphenoxy)benzaldehyde
	21126-5	4-Phenoxybenzaldehyde
	12371-4	4-Benzyloxybenzaldehyde
	16361-9	4-Benzyloxy-3-methoxybenzaldehyde
	16395-3	3-Benzyloxy-4-methoxybenzaldehyde
25	34603-9	3-Methoxy-4-(4-nitrobenzyloxy)benzaldehyde
	D3600-3	3,4-Dibenzyloxybenzaldehyde
	N10-9	1-Naphthaldehyde
	N20-6	2-Naphthaldehyde
	15134-3	2-Methoxy-1-naphthaldehyde
30	10324-1	4-Methoxy-1-naphthaldehyde
	H4535-3	2-Hydroxy-1-naphthaldehyde
	27208-6	4-Dimethylamino-1-naphthaldehyde
	38201-9	2,3-Naphthalendicarboxaldehyde
	15014-2	2-Fluorenenecarboxaldehyde
35	27868-8	9-Anthraldehyde

- 112 -

M2965-7 10-Methylanthracene-9-carboxaldehyde
15211-0 10-Chloro-9-anthraldehyde
P1160-3 Phenanthrene-9-carboxaldehyde
14403-7 1-Pyrenecarboxaldehyde

5 Aliphatic aldehydes

25254-9 Formaldehylde
11007-8 Acetaldehyde
P5145-1 Propionaldehyde
24078-8 Isobutyraldehyde
10 T7150-1 Trimethylacetaldehyde
B10328-4 Butyraldehyde
M3347-6 2-Methylbutyraldehyde
11009-4 2-Ethylbutyraldehyde
14645-5 Isovaleraldehyde
15 35990-4 3,3-Dimethylbutyraldehyde
11013-2 Valeraldehyde
25856-3 2-Methylvaleraldehyde
D19050-0 2,4-Dimethylvaleraldehyde
11560-6 Hexanal
20 E2910-9 2-Ethylhexanal
30355-0 3,5,5-Trimethylhexanal
H212-0 Heptaldehyde
O560-8 Octyl aldehyde
N3080-3 Nonyl aldehyde
25 12577-6 Decyl aldehyde
U220-2 Undecylic aldehyde
M8675-8 2-Methylundecanal
D22200-3 Dodecyl aldehyde
26923-9 Tridecanal
30 T1000-6 Tetradecy aldehyde
11022-1 Acrolein
13303-5 Methacrolein
25614-5 2-Ethylacrolein
25613-7 2-Butylacrolein

	13298-5	Crotonaldehyde
	19261-9	<i>trans</i> -2-Methyl-2-butenal
	29468-3	2-Ethyl- <i>trans</i> -2-butenal
	30407-7	3-Methyl-2-butenal
5	26925-5	<i>trans</i> -2-pentenal
	29466-7	2-Methyl-2-pentenal
	29097-1	2,2-Dimethyl-4-pentenal
	13265-9	<i>trans</i> -2-Hexenal
	25176-3	<i>trans</i> -2-Heptenal
10	30796-3	2,6-Dimethyl-5-heptenal
	26995-6	<i>trans</i> -2-Octenal
	34364-1	(R)-(+)-Citronellal
	37375-3	(S)-(-)-Citronellal
	25565-3	<i>trans</i> -2-Nonenal
15	37562-4	<i>cis</i> -4-Decenal
	36733-8	<i>trans</i> -4-Decenal
	13227-6	Undecylenic aldehyde
	24911-4	<i>dis</i> -9-hexadecenal
	27221-3	Cyclopropanecarboxaldehyde
20	10846-4	Cyclohexanecarboxaldehyde
	10933-9	Cyclooctanecarboxaldehyde
	30441-7	3-Cyclohexylpropionaldehyde
	T1220-3	Tetrahydrobenzaldehyde
	21829-4	(S)-(-)-Perillaldehyde
25	26467-9	2,6,6-Trimethyl-1-cyclohexene-1-acetaldehyde
	10937-1	5-Norbornen-2-carboxaldehyde
	21824-3	(1R)-(-)-Myrtenal
	37531-4	Glyoxal-1,1-dimethyl acetal
	21877-4	7-Methoxy-3,7-dimethyloctanal
30	23254-8	3-Ethoxymethacrolein
	27525-5	2,5-Dimethoxy-3-tetrahydrofurancarboxaldehyde
	26918-2	2,2-Dimethyl-3-hydroxypropionaldehyde
	G480-2	DL-Glyceraldehyde
	G478-0	D-Glyceraldehyde
35	21665-8	L-Glyceraldehyde

- 114 -

	34140-1	3-(Methylthio)propionaldehyde
	30583-9	3-(Dimethylamino)acrolein
	36549-9	3-(Dimethylamino)-2-methyl-2-propenal
	17733-4	Pyruvic aldehyde
5	27706-1	(S)-(-)-2-(Methoxymethyl)-1-pyrrolidinecarboxaldehyde
	29211-7	2-Methoxy-1-pyrrolidinecarboxaldehyde
	29210-9	2-Methoxy-1-piperidinecarboxaldehyde

EXAMPLE 123

10 **Synthesis of libraries from oligomeric compounds utilizing aryl acid halides. Use of benzoyl chloride Aldrich; 3-methylbenzoyl chloride Aldrich-T3,550-5; 3-methoxybenzoyl chloride, Aldrich-12,965-8; and 3-nitrobenzoyl chloride Aldrich-N1,084-5, as illustrative letters**

15 Preparation of combinatorial libraries as per Example 121 using benzoic acids and benzoic acid derivatives is effected in place of benzyl aldehydes described above using the general procedure of Examples 63 and 65.

20 Using the above, libraries are prepared from oligomeric compounds that are diveratized with one, two, three, four or more of the following acid halides available from Aldrich Chemical Company, Inc., Milwaukee, WI. The Aldrich catalog number is given in the right hand column and the compound name is given in the left hand column:

25	10663-1	p-Toluoyl chloride
	30253-8	3-Cyanobenzoyl chloride
	13096-6	(+/-)-2-Chloro-2-phenylacetyl chloride
	26366-4	3-(Chloromethyl)benzoyl chloride
	27078-4	4-(Chloromethyl)benzoyl chloride
30	24947-5	4-(Trifluoromethyl)benzoyl chloride
	19394-1	4-Chlorophenoxyacetyl chloride
	24948-3	2-(Trifluoromethyl)benzoyl chloride
	19394-1	4-Chlorophenoxyacetyl chloride

- 115 -

	24948-3	2-(Trifluoromethyl)benzoyl chloride
	10663-1	<i>p</i> -Toluoyl chloride
	25027-9	3-(Trifluoromethyl)benzoyl chloride
	S67828-7	2-(2,4,5-Trichlorophenoxy)acetyl chloride
5	12201-7	<i>o</i> -Toluoyl chloride
	40248-6	4-(Trifluoromethoxy)benzoyl chloride
	37502-0	3-(Dichloromethyl)benzoyl chloride
	12225-4	<i>m</i> -Toluoyl chloride
	12482-6	4-Cyanobenzoyl chloride
10	P1675-3	Phenylacetyl chloride
	S88415-4	2-(Phenylthio)propionyl chloride
	15862-3	Phenoxyacetyl chloride
	36475-4	<i>trans</i> -4-Nitrocinnamoyl chloride
	28882-9	4-Ethoxybenzoyl chloride
15	23024-3	<i>m</i> -Anisoyl chloride
	S67595-4	2,3-Dibromo-3-phenylpropionyl chloride
	30101-9	Benzyloxyacetyl chloride
	25470-3	<i>o</i> -Anisoyl chloride
	C8110-1	Cinnamoyl chloride
20	31693-8	3-Methoxyphenylacetyl chloride
	A8847-6	<i>p</i> -Anisoyl chloride
	16519-0	Acetylsalicyloyl chloride
	36569-6	4-Methoxyphenylacetyl chloride
	24944-0	Hydrocinnamoyl chloride
25	26528-4	3,5-Bis(trifluoromethyl)benzoyl chloride
	28350-94	Ethylbenzoyl chloride
	S40503-5	2-Phenoxypropionyl chloride
	33304-2	2,5-Bis(trifluoromethyl)benzoyl chloride
	S62043-2	<i>p</i> -Tolylacetyl chloride
30	16171-3	3,5-Dimethoxybenzoyl chloride
	42339-4	(R)-(-)- <i>A</i> -Methoxy- <i>A</i> -(trifluoromethyl)- phenylacetyl chloride
	26480-6	2,5-Dimethoxyphenylacetyl chloride
	25804-0	3,4-Dimethoxybenzoyl chloride
35	T6980-9	3,4,5-Trimethoxybenzoyl chloride

- 116 -

	26242-0	2,6-Dimethoxybenzoyl chloride
	13430-9	trans-2-Phenyl-1-cyclopropanecarbonyl chloride
	S62264-8	5-(Dimethylsulfamoyl)-2-methoxybenzoyl chloride
5	37383-4	2,4-Dimethoxybenzoyl chloride
	A1740-4	o-Acetylmandelic chloride
	24945-9	4-Phenyl-1,2,3,4-tetrachloro-1,3-butadiene-1-carbonyl chloride
	36848-2	trans-3-(trifluoromethyl)cinnamoyl chloride
10	15712-0	4-tert-butylbenzoyl chloride
	S42860-4	2-Phenylbutyryl chloride
	22203-8	4-Butylbenzoyl chloride
	23747-7	3,4-Dimethoxyphenylacetyl chloride
	22204-6	4-Butoxybenzoyl chloride
15	S65659-3	2-(4-Chlorobenzoyl)benzoyl chloride
	22214-3	4-Pentylbenzoyl chloride
	C3928-8	2-Chloro-2,2-diphenylacetyl chloride
	S43639-9	4(4-Nitrophenylazo)benzoyl chloride
	33158-9	Diphenylacetyl chloride
20	S80926-8	4-(Phenylazo)benzoyl chloride
	S61661-3	2-Diphenylacetyl chloride
	16114-4	4-Biphenylcarbonyl chloride
	22209-7	4-Hexylbenzoyl chloride
	22205-4	4-Heptyloxybenzoyl chloride
25	22211-9	4-Hexyloxybenzoyl chloride
	22206-2	4-Heptyloxybenzoyl chloride

EXAMPLE 124**Loading of solid support, General Procedure****Method 1: Succinylation and activation of aminodiol monomer**

- 30 An Fmoc protected mono-DMT aminodiol monomer (36 mmol) was dissolved in 400 ml CH₂Cl₂, and triethylamine (45 mmol, 6.25 ml) and succinic anhydride (40 mmol, 4.0 g) added. A catalytic amount of DMAP was added (5 mol %) and the solution

- 117 -

stirred at room temperature overnight. TLC showed all the starting material was converted to a more polar spot. To the solution was then added pentafluorophenyl trifluoroacetate (6.9 ml, 40 mmol). The solution was stirred 1 hr, and the polar material was converted to a non-polar spot on TLC. Toluene (150 ml) was added, and the solvents evaporated. The oily residue was loaded on a flash chromatography column, and eluted with ethyl acetate and hexanes to give the product (33 mmol, 93 %).

10 Method 2: Derivatization of TentaGel

TentaGel-NH₂ (50 g, 11 mmol NH₂) was swelled in 150 ml CH₂Cl₂, then washed with 5 % diisopropylethylamine in CH₂Cl₂, followed by CH₂Cl₂. The resin was sucked dry, and a solution of monomer succinate PFP ester (17 mmol) in 120 ml CH₂Cl₂ and triethylamine (3.5 ml, 25 mmol) were added. The solution was agitated on a wrist shaker for 6 hours, filtered and rinsed with CH₂Cl₂, and a second portion of PFP ester (16 mmol) and triethylamine (3.5 ml) added. The solid support was shaken overnight, filtered and resuspended in 60 ml CH₂Cl₂, 60 ml pyridine and 10 ml acetic anhydride added. After 1 hr, the resin was filtered and washed with CH₂Cl₂, pyridine/CH₂Cl₂/Methanol (1:8:1), and diethyl ether. The resin was dried under high vacuum. The loading was determined by DMT cation to be 0.14-0.20 mmol/g.

25 EXAMPLE 125

Derivatization of scaffolds on solid support. General procedure for acylation of scaffold monomers with carboxylic acids.

An aminodiol monomer on a solid support (0.5 g Tentagel, 0.15 mmol/g) is placed in a shaker flask and purged with argon for 15 minutes. The support is pre-swelled in CH₂Cl₂ (60 min.) then washed with DMF (6 ml). The Fmoc protecting group,

- 118 -

if present, is removed by addition of piperidine/DMF (10%, 6 mL, ~80 eq.), followed by agitation of the reaction mixture for 15 minutes. The support was washed with DMF (6 ml X 5). A solution of a carboxylic acid (0.4 M, ~5 eq.) and DIEA (0.8

5 M, ~10 eq.) in DMF was added, followed by a solution of activator, BOP or HATU in DMF (0.4 M, 5 eq). The reaction mixture was agitated (30 min.) and then washed with DMF (6 ml X 3) and CH₂Cl₂ (6 ml X 3). Several other activating agents can be used. These include carbodiimides such as

10 dicyclohexyl carbodiimide (DCC), diisopropyl carbodiimide (DIC), dimethylaminopropyl ethyl carbodiimide hydrochloride (EDC), EEDQ and IIDQ, carbonyl diimidazole, with or without the addition of additives such as dimethylaminopyridine, N-hydroxybenzotriazole, N-hydroxy-7-azabenzotriazole, and

15 others known to those skilled in the art. Other useful coupling agents include substituted uronium salts and phosphonium salts such as HBTU, TBTU, BOP, PyBROP and other analogous reagents, and reagents for producing acyl fluorides, such as cyanuric fluoride. Representative examples

20 of each of these reagents can be found in Bodanszky, M. Principles of Peptide Synthesis, 2nd Ed. Springer-Verlag, Berlin, 1993.

Representative carboxylic acids that can be used to derivatize an amino group of an aminodiol of the invention

25 include:

2-oxovaleric acid

2-oxo-octanoic acid

2-oxo-2-(2-furyl)acetic acid

indole-3-pyruvic acid

30 2-nitrophenylpyruvic acid

2-furylthiopyruvic acid

methacrylic acid

2-methylpropionic acid (isobutyric acid)

cyanoacetic acid

35 methoxyacetic acid

- 3-methylthiopropionic acid
4-methylpentanoic acid
3-trimethylsilylpropionic acid sodium salt
N-BOC-5-aminovaleric acid]
- 5 3-(N,N-diethylamino)propionic acid hydrochloride
monomethyl glutarate
7-oxo-octanoic acid
neodecanoic acid
1,2,3-thiadiazole-4-carboxylic acid
- 10 3-amino-1,2,4-triazole-5-carboxylic acid
3-furoic acid
2-furoic acid
4-(S)-butyrolactone-4-carboxylic acid
1-methylpyrrole-2-carboxylic acid
- 15 4-methyl-2-phenyl-1,2,3-triazole-5-carboxylic acid
1-(3'-aminophenyl)-3-carboxy-5-pyrazolone
4-phenyl-5-trifluoromethyl-thiophene-2-carboxylic acid
3-(4-chlorophenylthio)thiophene-4-carboxylic acid
5-(2-pyridylthiomethyl)-2-furancarboxylic acid
- 20 indole-2-carboxylic acid
1-methylindole-2-carboxylic acid
7-benzyloxyindole-2-carboxylic acid
4-oxo-4,5,6,7-tetrahydrobenzo[b]furan-3-carboxylic acid
1-methylindene-2-carboxylic acid
- 25 4-chloro-3-sulfamoylbenzoic acid
2-hydroxybenzoic acid (salicylic acid)
a-mercapto-p-toluic acid
BOC 4-(aminomethyl)benzoic acid
BOC 4-(methylamino)benzoic acid
- 30 N-acetyl-4-aminobenzoic acid
4-isopropoxybenzoic acid
4-(2-[methylsulfonamido]ethoxy) benzoic acid
4-(1H-pyrrol-1-yl)benzoic acid
4-(3-methyl-5-oxo-2-pyrazolin-1-yl)benzoic acid
- 35 4'-(trifluoromethyl)-2-biphenylcarboxylic acid

- 120 -

- 2-(4-nitrophenylthio)benzoic acid
2-phenoxybenzoic acid
2-(4-chlorobenzoyl)benzoic acid
2-benzoylbenzoic acid
5 a-phenyl-o-toluic acid
a-(4-methoxy-1-naphthyl)-o-toluic acid
2-(1,2,3,4-tetrahydro-6-naphthylmethyl)benzoic acid
2-(2-pyridylcarbonyl)benzoic acid
4-(3-fluorobenzamido)benzoic acid
10 3'-carboxy-3-methylbenzanilide
2'-methoxyphthalanilic acid
3-benzyloxy-4-methoxybenzoic acid
N-(3-methyl-2-pyridyl)phthalamic acid
1-(4-carboxyphenyl)-3-(o-tolyl)urea
15 benzotriazole-5-carboxylic acid
5-benzimidazolecarboxylic acid
piperonylic acid
2,2,5,7-tetramethylindan-1-one-4-carboxylic acid
2-naphthoic acid
20 2-pyrazinecarboxylic acid
picolinic acid
nicotinic acid
isonicotinic acid
6-hydroxynicotinic acid
25 tetrahydropyran-4-carboxylic acid
N-BOC 4-piperidinecarboxylic acid (isonipecotinic acid)
1,3-dimethyl-6-uracilcarboxylic acid
2-pyrrolopyridine-5-carboxylic acid
2-phenoxy nicotinic acid
30 2-(4-methylphenoxy)pyridine-3-carboxylic acid
2-phenylimidazo(1,2-A)pyridine-6-carboxylic acid
3-isoquinolinecarboxylic acid
4-quinolinecarboxylic acid
4-hydroxy-7-trifluoromethyl-3-quinolinecarboxylic acid
35 2-p-tolylcinchoninic acid

- 121 -

- nalidixic acid
thiophene-2-acetic acid
5-(1-pyrrolidine)2-tetrazoleacetic acid
N-phthaloylglycine
5 indole-3-acetic acid
cyclopentylacetic acid
2-indanylacetic acid
pentafluorophenylacetic acid
2-(2-fluorophenyl)acetic acid
10 (S)-(+)-mandelic acid
(a,a,a-trifluoro-m-tolyl)acetic acid
R-(-)-a-methoxyphenylacetic acid
2-(4-chlorophenyl)-2,2-dimethylacetic acid
(R)-(-)-2-phenylbutyric acid
15 (3,4-dimethoxyphenyl)acetic acid]
4-(dimethylamino)phenylacetic acid
4-biphenylacetic acid
2-benzyloxyphenylacetic acid
1-(4-methylphenyl)-1-cyclopropanecarboxylic acid
20 benzilic acid
orotic acid
4-pyridylacetic acid hydrochloride
phenylacetic acid
2-(2-phenyl-1-cyclohexenyl)acetic acid
25 trans-3-furanacrylic acid
2-methyl-4-nitro-1-imidazolepropionic acid
5-phenyl-2-pyrrolepropionic acid
2-(4-chlorophenoxy)acetic acid
phenoxyacetic acid
30 N-phenylglycine
2,5-difluorocinnamic acid
3-(4-iodophenyl)propionic acid
N-(2,4-dinitrophenyl)-L-alanine
3-(3-hydroxyphenyl)propionic acid
35 p-toluenesulfonylacetic acid

- 122 -

- BOC-3-(p-aminophenyl)propionic acid
4-cyanocinnamic acid
4-methoxycinnamic acid
2-(p-chlorophenoxy)-2-methylpropionic acid
5 p-(p-nitrobenzyloxy)cinnamic acid
2-naphthoxyacetic acid
3,3-diphenylpropionic acid
3,5,6-trichloro-2-pyridoxyacetic acid
4-pyridylthioacetic acid
10 5-trifluoromethyl-2-pyridylthioacetic acid
1-piperidinepropionic acid
4-oxo-4-(1-pyrrolidine)butyric acid
3-(4-fluorobenzoyl)propionic acid
3-benzoylpropionic acid
15 N-methylhippuric acid
4-oxo-4-(2-trifluoromethylphenyl)butyric acid
3-(4-methoxybenzoyl)propionic acid
3-(4-methylsulfonylbenzoyl)propionic acid
4-(4-acetylphenyl)butyric acid
20 4-oxo-4-(2-naphthyl)butyric acid
dansyl glycine
3-(4-methoxy-1-naphthoyl)propionic acid
(-)-O,O'-dibenzoyl-L-tartaric acid mono(dimethylamide)
1-(3-carboxy-1-oxopropyl)-1,2,3,4-tetrahydroquinoline
25 1,2-dihydro-2-methyl-1-oxo-3-isoquinolinebutyric acid
2-(2-carboxyethylthio)-3,5,6-trimethyl-1,4-benzoquinone
(+)-biotin
4-(2-mercaptobenzothiazolyl)butyric acid
Z-styrenesulfonylacetic acid
30 4-(p-chlorophenoxy)butyric acid
3-(benzylthio)propionic acid
N-(2-pyridyl)succinamic acid
Cbz-glycine
2-[(2-phenoxyethyl)thio]acetic acid
35 2-(benzenesulfonyl)ethylthioacetic acid

- 123 -

2'-carbamoylglutaranilic acid.

EXAMPLE 126

Acylation of scaffold monomers with acyl chlorides, anhydrides, and activated carboxylic acid derivatives.

5 An aminodiol monomer on a solid support (0.5 g Tentagel, 0.15 mmol/g) was placed in a shaker flask and purged with argon (15 min.). The support was pre-swelled in CH_2Cl_2 (60 min.) then washed with DMF (6 ml). The Fmoc protecting group, if present, is removed by addition of piperidine/DMF (10%, 6
10 mL, ~80 eq.), followed by agitation of the reaction mixture for 15 minutes. The support was washed with DMF (6 ml X 5). A solution of py/DMF (10%, 3 mL, ~50 eq.), acid chloride/DMF (0.12 M, 3 mL, ~5 eq.) was added, and the reaction mixture agitated for 30 minutes. A solution of acid anhydride or
15 other activated derivative in DMF could be used instead of acid chloride. The support was then washed with DMF (6 ml X 3) and CH_2Cl_2 (6 ml X 3). Examples of reagents which can be substituted for acid chlorides are: acid anhydrides and mixed anhydrides, imidazolides, or active esters such as N-hydroxy
20 succinimide esters and other O-acyl hydroxylamine derivatives, acyl azides, 4-nitrophenol esters, pentachlorophenyl or pentafluorophenyl esters and other active aryl and vinyl esters. Representative examples of each of these reagents can be found in Bodanszky, M.
25 Principles of Peptide Synthesis, 2nd Ed. Springer-Verlag, Berlin, 1993.

EXAMPLE 127

Sulfonylation of scaffold monomers:

30 An aminodiol monomer on a solid support (0.5 g Tentagel, 0.15 mmol/g) was placed in a shaker flask and purged with argon (15 min.). The support was pre-swelled in CH_2Cl_2 (60 min.) then washed with DMF (6 ml). The Fmoc protecting group,

- 124 -

if present, is removed by addition of piperidine/DMF (10%, 6 mL, ~80 eq.), followed by agitation of the reaction mixture for 15 minutes. The support was washed with DMF (6 ml X 5). A solution of TEA/CH₃CN (10%, 3 mL, ~30 eq.) was added
5 followed by the sulfonyl chloride in CH₃CN/Pyridine (0.12 M, 3 mL, ~5 eq.), and the reaction mixture agitated for 30 minutes. The support was washed with pyridine (6 ml X 3), DMF (6 ml X 3) and CH₂Cl₂ (6 ml X 3).

Representative sulfonyl chlorides suitable for use in
10 the invention include:

methanesulfonyl chloride
dimethylsulfamoyl chloride
1--butanesulfonyl chloride
3,5-dimethylisoxazole-4-sulfonyl chloride
15 5-(4-chlorobenzamidomethyl) thiophene-2-sulfonyl chloride
benzenesulfonyl chloride
o-carbomethoxybenzenesulfonyl chloride
N-acetylsulfanilyl chloride
2,5-dimethoxybenzenesulfonyl chloride
20 3-(chlorosulfonyl)benzoic acid
2-dibenzofuransulfonyl chloride
2-naphthalenesulfonyl chloride
8-quinolinesulfonyl chloride
phenylmethanesulfonyl chloride
25 (+/-)-10-camphorsulfonyl chloride
N-acetylsulfanilyl chloride
3,5-dichlorobenzene sulfonyl chloride
dansyl chloride.

EXAMPLE 128

30 Alkylation of scaffold monomers

An aminodiol monomer on a solid support (0.5 g Tentagel, 0.15 mmol/g) was placed in a shaker flask and purged with argon (15 min.). The support was pre-swelled in CH₂Cl₂ (60 min.) then washed with DMF (6 ml). The Fmoc protecting group,

- 125 -

if present, is removed by addition of piperidine/DMF (10%, 6 mL, ~80 eq.), followed by agitation of the reaction mixture for 15 minutes. The support was washed with DMF (6 ml X 5). A solution of DIEA in DMF (10%, 3 mL, ~50 eq.), alkylating reagent (Alkyl halide, mesylate, tosylate or triflate) in DMF (0.2 M, 3 mL, ~5 eq.) is added, and the reaction mixture is agitated for 120 min. The support is then washed with DMF (6 ml X 3) and CH₂Cl₂ (6 ml X 3).

EXAMPLE 129**10 Synthesis of Ureas on Solid Support**

An aminodiol monomer on a solid support (0.5 g Tentagel, 0.15 mmol/g) was placed in a shaker flask and purged with argon (15 min.). The support was pre-swelled in CH₂Cl₂ (60 min.) then washed with DMF (6 ml). The Fmoc protecting group, if present, is removed by addition of piperidine/DMF (10%, 6 mL, ~80 eq.), followed by agitation of the reaction mixture for 15 minutes. The support was washed with DMF (6 ml X 5). A solution of isocyanate in DMF (2.0 ml, 0.2M) is added, followed by DMAP in DMF (2.0 ml, 0.02M) and the resulting mixture agitated (120 min.). The support is then washed with DMF (6 ml X 3) and CH₂Cl₂ (6 ml X 3).

Representative isocyanates suitable for use in the invention include:

- benzoyl isocyanate
- 25 4-chlorobenzenesulfonyl isocyanate
- benzenesulfonyl isocyanate
- 3-nitrophenyl isocyanate
- 4-acetylphenyl isocyanate
- 2,5-dimethylphenyl isocyanate
- 30 ethyl 4-isocyanatobenzoate
- 2-phenylphenyl isocyanate
- 4-phenoxyphenyl isocyanate
- 1-naphthyl isocyanate
- R-(+)-α-methylbenzyl isocyanate

- 126 -

Methyl Isocyanate

Isopropyl Isocyanate

Cyclohexyl isocyanate

Allyl Isocyanate

5 Chloromethyl Isocyanate

Chloropropyl Isocyanate

Ethyl Isocyanatoacetate

4-(Chloromethyl) phenyl isocyanate

3-Chlorophenyl Isocyanate

10 4-Fluorophenyl Isocyanate

Ethyl 4-Isocyanatobenzoate

3,5-Dichlorophenyl Isocyanate

EXAMPLE 130**Quality Control:**

15 An aliquot (5 mg) of a scaffold attached to a solid support that has been derivatized is placed in a 1 mL disposable syringe fitted with glass wool plug. TCA/CH₂Cl₂ (3%, 0.4 ml X 5) is drawn in, and the reaction mixture is agitated manually to remove the DMT. The support is washed

20 with CH₂Cl₂ (6 ml X 3) and pyridine-CH₃CN (6 ml X 3). DMT-deoxythymidine H-phosphonate in pyridine-CH₃CN (0.1 M, 0.4 mL, ~50 eq.) is added, followed by a solution of adamantoyl chloride in pyridine-CH₃CN, and the reaction mixture is agitated manually for 3 minutes. The support is washed with

25 pyridine-CH₃CN (1 ml X 3) and THF (1 ml X 3). Oxidation of the H-phosphonate diester is achieved by adding a solution of I₂ in THF (0.2 M, 0.4 mL, ~100 eq.) and H₂O/TEA/THF (1:1:8, 0.4 mL), and agitating periodically. The support is washed with THF (1 mL X 3), pyridine-CH₃CN (1 mL X 5), DMF (1 mL X

30 3), CH₂Cl₂ (1 mL X 3), and MeOH (1 mL X 3). NH₄OH (28-30%, 0.5 mL) is added to cleave the product from the support. After 1-2 hours, the ammonia solution is collected in vials and subjected to HPLC and mass spectrometric analyses.

EXAMPLE 131

(S)-2-*t*-Butoxycarbonyl-1,2,3,4-tetrahydro-7-hydroxyisoquinoline-3-carboxylic Acid.

The title compound was made by the method of
5 Verschueren, K., Thoth, G., Tourwé, D., Lebl, M., Van Binst, G., Hruby, V. *Synthesis*, 1992, 5, 458-4602.

EXAMPLE 132

(S)-2-*t*-Butoxycarbonyl-1,2,3,4-tetrahydro-3-hydroxymethyl-7-hydroxyisoquinoline.

10 (S)-2-*t*-Butoxycarbonyl-1,2,3,4-tetrahydro-7-hydroxyisoquinoline-3-carboxylic acid (100 mmol) is dissolved in THF (1 L). Borane methyl sulfide complex (200 mmol) is added carefully, and the solution heated at reflux for 1 hour. The solution is cooled, quenched with MeOH, and 1 N
15 HCl to pH 2. The solution is neutralized to pH 7-8 with NaHCO₃ and the solvent evaporated. The oily residue is diluted with water, extracted with EtOAC, washed with NaHCO₃ and brine, then dried with MgSO₄ and evaporated to give the title compound.

20 EXAMPLE 133

(S)-2-*t*-Butoxycarbonyl-3-hydroxymethyl-7-hydroxyoctahydroisoquinoline

To a solution of *t*-butyl alcohol (9.1 g, 123 mmol), 50 ml ether, and ammonia (150 mL) is added (S)-2-*t*-
25 butoxycarbonyl-1,2,3,4-tetrahydro-3-hydroxymethyl-7-hydroxyisoquinoline (5 mmol). The solution is brought to reflux, and lithium shot (0.51 g, 82 mmol) is added over 30 minutes. The solution is allowed to reflux for 3 hours, and solid NH₄Cl added until the color disappears. The ammonia is
30 allowed to evaporate overnight, ice-water is added, and the organic layer separated. The aqueous layer is extracted with chloroform and the combined organic phases are washed with

- 128 -

brine, dried and evaporated. The crude product is redissolved in methanol, 5 % Pt/C added and the solution shaken under 3 atm H₂ for 6 hours to complete the reduction. The catalyst is filtered and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

EXAMPLE 134**(S)-2-FMOC-3-hydroxymethyl-7-hydroxyoctahydroisoquinoline**

The Boc protected material from the previous example is dissolved in ethyl acetate and 1 N HCl in ethyl acetate is added. The solution is stirred until all the starting material is consumed. The solvent is evaporated and the residue redissolved in 2M NaHCO₃ (25 ml) and dioxane (25 ml). FMOC-Cl is added and the solution stirred for 6 hours. The solvent is evaporated, and the product is extracted with ethyl acetate, washed with NaHCO₃, and then brine, dried with MgSO₄, and evaporated. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

EXAMPLE 135**(S)-2-FMOC-3-dimethoxytrityloxymethyl-7-hydroxyoctahydroisoquinoline**

The product from the previous example is dissolved in pyridine, cooled to 0 °C and 1.2 eq DMT-Cl added. The solution is stirred overnight at 0 °C, quenched with methanol, and the solvent evaporated. The residue is redissolved in EtOAc, washed with 5 % citric acid, NaHCO₃, and brine, and evaporated. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

EXAMPLE 136

1-BOC-3-hydroxymethylpyrrole

The title compound is prepared according to the method of Davies, H. M. L., Matasi, J. J., Ahmed, G. J. *Org.Chem.* 1996, 61, 2305.

5 **EXAMPLE 137****1-BOC-3-O-TBDMS-hydroxymethylpyrrole**

1-BOC-3-hydroxymethylpyrrole (2.5 eq) and imidazole (2.5 eq) are dissolved in DMF and TBDMS-Cl (1.2 eq) is added. The solution is stirred overnight, quenched with water, and
10 extracted with EtOAc. The extracts are washed with water, NaHCO₃, and brine, and dried with MgSO₄. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

EXAMPLE 13815 **Methyl-3-O-TBDMS-6-(O-TBDMS)-oxymethyl-8-BOC-8-azabicyclo[3.2.1.]octa-2,6-diene-2-carboxylate**

A solution of methyl 2-diazoacetoacetate (25 mmol) and triethylamine (30 mmol) in CH₂Cl₂ is cooled to 0 °C, and TBDMS-triflate (27 mmol) is added. The solution is stirred
20 at 0 °C for 30 minutes, diluted with hexane, and washed with 5% NaHCO₃. The extracts are dried and evaporated, and the crude enol silyl ether is used directly. The enol ether is dissolved in hexane (50 mL), and added slowly to a solution of 1-BOC-3-O-TBDMS-hydroxymethylpyrrole (20 mmol) and rhodium
25 (II) hexanoate in refluxing hexane (50 mL) under an atmosphere of argon. The solution is heated a further 12 hours and evaporated. The product is purified by flash chromatography. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

30 **EXAMPLE 139****Methyl-3-O-TBDMS-6-(O-TBDMS)-oxymethyl-8-BOC-8-**

- 130 -

azabicyclo[3.2.1.]octa-2-ene-2-carboxylate

The diene product of the previous example and tris(triphenylphosphine) rhodium chloride (5 mol%) are dissolved in ethanol, and shaken under 45 psi H₂ for 12 hours.

- 5 The solvent is evaporated and the resulting residue is purified by silica gel flash column chromatography to give the title compound.

EXAMPLE 140**Methyl-3-oxo-6-hydroxymethyl-8-BOC-8-**

- 10 **azabicyclo[3.2.1.]octane-2-carboxylate**

- The bis silyl ether of the previous example is dissolved in THF, and a solution of tetrabutylammonium fluoride in THF is added (3 eq). The solution is stirred at room temperature until the starting material is consumed. The solvent is
15 evaporated, the residue redissolved in ether and washed with dilute NaHCO₃, brine, and dried with MgSO₄. The crude material is filtered through silica gel to give the title compound.

EXAMPLE 141

- 20 **3-Oxo-6-hydroxymethyl-8-BOC-8-azabicyclo[3.2.1.]octane**

- The keto ester of the previous example is dissolved in DMSO, and 1 eq NaCl and 1 eq H₂O added. The solution is heated to 160 °C for 1 hour in an open vessel, cooled to room temperature and diluted with water. The suspension is
25 extracted with EtOAc, washed with 5 % NaHCO₃, brine, dried over MgSO₄, and evaporated. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

EXAMPLE 142

- 30 **3-Oxo-6-O-DMT-oxymethyl-8-FMOC-8-azabicyclo[3.2.1.]octane**

3-Oxo-6-hydroxymethyl-8-BOC-8-azabicyclo[3.2.1.]octane

- 131 -

is dissolved in ethyl acetate and 1 N HCl in ethyl acetate is added. The solution is stirred until all the starting material is consumed. The solvent is evaporated and the residue redissolved in 2M NaHCO₃ (25mL) and dioxane (25 mL).

5 FMOC-Cl is added and the solution stirred for 6 hours. The solvent is evaporated, and the product extracted with ethyl acetate, washed with NaHCO₃, brine, dried over MgSO₄, and evaporated. The crude material is dissolved in pyridine, cooled to 0 °C and 1.2 eq DMT-Cl added. The solution is
10 stirred overnight at 0 °C, quenched with methanol, and the solvent evaporated. The residue is redissolved in EtOAc, washed with 5 % citric acid, NaHCO₃, brine and evaporated. The resulting residue is purified by silica gel flash column chromatography to give the title compound.

15 **EXAMPLE 143**

3-Hydroxy-6-O-DMT-oxymethyl-8-FMOC-8-azabicyclo[3.2.1.]octane

The product from the previous example is dissolved in THF and 1 eq of a solution of LiBH₄ is added. The solution is stirred until the starting material was consumed. The
20 reaction is carefully quenched to pH 4 (wet litmus), and neutralized with NaHCO₃. The solvent is evaporated, the residue extracted with ethyl acetate, washed with NaHCO₃, brine, dried and evaporated. The product is purified by flash chromatography.

25 **EXAMPLE 144**

N-Methyl-2,6-diacetoxy-9-azabicyclo[3.3.1]nonane

The title compound is made by the method of Clousdale, I. S., Kluge, A. F., McClure, N. L. *J. Org. Chem.* 1982, 47, 919.

30 **EXAMPLE 145**

2,6-Diacetoxy-9-azabicyclo[3.3.1]nonane

- 132 -

The material from the previous example (10 mmol) is dissolved in 100 ml 1,2-dichloroethane, cooled to 0 °C and 1-chloroethyl chloroformate (11 mmol) is added. The solution is warmed to reflux temperature after 15 minutes, and heated until the starting material is consumed. The solvent is then removed, and the residue redissolved in anhydrous methanol, and warmed to 50 °C for 1 hour. The solvent is removed to give the title compound as the HCl salt.

EXAMPLE 146**2,6-Dihydroxy-9-FMOC-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) is dissolved in 50 ml 1:1 aqueous THF, and 50 mmol LiOH added. The biphasic solution is stirred at room temperature for 6 hours. A solution of FMOC-Cl (11 mmol) in THF is then added dropwise at room temperature over 15 minutes. After 1 hour the solution is acidified with 10% HCl to pH 2, and extracted with EtOAc, washed with water, brine and dried with MgSO₄. The product is purified by flash chromatography.

EXAMPLE 147**2-O-DMT-6-hydroxy-9-FMOC-9-azabicyclo[3.3.1]nonane**

The diol from the previous example (10 mmol) is dissolved in CH₂Cl₂ and pyridine (9:1), cooled to 0 °C and DMT-Cl (10 mmol) is added in portions (1.0 eq). The solution is stirred overnight at 0 °C. The reaction is quenched with MeOH, evaporated, and excess solvent is removed by azeotropic distillation with toluene. The residue is redissolved in EtOAc, washed with 5% citric acid, NaHCO₃, brine and dried with MgSO₄. The product is purified by flash chromatography using a gradient of MeOH in CH₂Cl₂. The product is obtained in 50% yield, along with 25% ditritylated material which is conserved and recycled, and 25% unreacted starting material.

EXAMPLE 148**Dimethyl 3-O-TBDMS-glutarate**

Dimethyl 3-hydroxyglutarate (34 mmol) and imidazole (85 mmol) are dissolved in 150 ml DMF, and TBDMS-Cl added (40 mmol). The solution is stirred overnight, quenched with water (400 ml), extracted with EtOAc, washed with NaHCO₃, brine and dried with MgSO₄ to give the title compound.

EXAMPLE 149**3-O-TBDMS-1,5-pentanediol**

The material from the previous example is dissolved in dry THF and 35 ml of 2M LiBH₄ in THF added. The solution is heated to reflux for 1.5 hrs, cooled to room temperature and quenched carefully with 10% HCl to pH 2, neutralized with NaHCO₃ and evaporated. The residue is extracted with EtOAc, washed with NaHCO₃, brine and dried with MgSO₄ to give the title compound.

EXAMPLE 150**N-Methyl-3-O-TBDMS-7-keto-9-azabicyclo[3.3.1]nonane**

To a stirred solution of oxalyl chloride (10 mmol) in 20 ml THF at -78 °C is added 11 mmol dimethylsulfoxide. The solution is warmed to -35 °C for 5 minutes and recooled to -78 °C. A solution of 3-O-TBDMS-1,5-pentanediol (5 mmol) in 10 ml THF is added, and the solution warmed to 0 °C. After 15 minutes, triethylamine (40 mmol) is added and the solution stirred at room temperature to effect complete conversion to the dialdehyde. The solvent is evaporated under reduced pressure, and the crude dialdehyde is redissolved in 40 ml citrate/phosphate buffer (pH 5.5) to which is added 15 mmol methylamine hydrochloride in 50 ml water, followed by the dropwise addition of 12 mmol acetone dicarboxylate in 80 ml water. The solution is stirred 24 hrs, 5 ml concentrated HCl added, and the solution heated to 90 °C for 1 hr to complete

the decarboxylation. Sodium hydroxide is added to pH 12, and the solution extracted with CH_2Cl_2 , dried over MgSO_4 and purified by chromatography on alumina to yield the product.

EXAMPLE 151**5 N-FMOC-3-O-TBDMS-7-keto-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) is dissolved in 100 ml 1,2-dichloroethane, cooled to 0 °C and 1-chloroethyl chloroformate (22 mmol) is added. The solution is warmed to reflux temperature after 15 minutes, and heated
10 until the starting material is consumed. The solvent is then removed, and the residue redissolved in anhydrous methanol, and warmed to 50 °C for 1 hour. The solvent is removed and the residue redissolved in 25 ml THF and 25 ml saturated NaHCO_3 . A solution of FMOC-Cl (11 mmol) in THF is then added
15 dropwise at room temperature over 15 minutes. After 1 hr the solution is acidified with 10% HCl to pH 2, and extracted with EtOAc, washed with water, brine and dried with MgSO_4 . The product is purified by flash chromatography.

EXAMPLE 152**20 N-FMOC-3-O-TBDMS-7-hydroxy-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) in dissolved in dry THF and 6 ml of 2 M LiBH_4 in THF added. The solution is stirred at room temperature and quenched carefully with 10% HCl to pH 2, neutralized with NaHCO_3 and
25 evaporated. The residue is extracted with EtOAc, washed with NaHCO_3 , brine and dried with MgSO_4 .

EXAMPLE 153**N-FMOC-3-O-TBDMS-7-O-DMT-9-azabicyclo[3.3.1]nonane**

The alcohol from the previous example (10 mmol) is
30 dissolved in CH_2Cl_2 and pyridine (9:1), cooled to 0 °C and DMT-Cl (11 mmol) added in portions (1.0 eq). The solution is

- 135 -

stirred overnight at room temperature. The reaction is quenched with MeOH, evaporated, and excess solvent removed by azeotropic distillation with toluene,. The residue is redissolved in EtOAc, washed with 5% citric acid, NaHCO₃,
5 brine and dried with MgSO₄. The product is purified by silica gel flash chromatography with a gradient of MeOH in CH₂Cl₂.

EXAMPLE 154**N-FMOC-3-hydroxy-7-O-DMT-9-azabicyclo[3.3.1]nonane**

The product of the previous example is added to a
10 solution of triethylamine trihydrofluoride (100 mmol) and triethylamine (50 mmol) in 50 ml of THF. The solution is stirred until the starting material is completely consumed. The reaction is stopped by the addition of saturated NaHCO₃, followed by evaporation of the solvent. The residue is
15 extracted with EtOAc, washed with NaHCO₃, brine and dried with MgSO₄. The product is purified by flash chromatography.

EXAMPLE 155**N-Methyl-2,6-diacetoxy-9-azabicyclo[3.3.1]nonane**

The title compound is made by the method of Clousdale, I.
20 S., Kluge, A. F., McClure, N. L. *J. Org. Chem.* **1982**, 47, 919.

EXAMPLE 156**2,6-Diacetoxy-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) is dissolved in 100 ml 1,2-dichloroethane, cooled to 0 °C and 1-
25 chloroethyl chloroformate (11 mmol) is added. The solution is warmed to reflux temperature after 15 minutes, and heated until the starting material is consumed. The solvent is then removed, and the residue redissolved in anhydrous methanol, and warmed to 50 °C for 1 hour. The solvent is removed to
30 give the title compound as the HCl salt.

- 136 -

EXAMPLE 157**2,6-Dihydroxy-9-FMOC-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) is dissolved in 50 ml 1:1 aqueous THF, and 50 mmol LiOH added. The biphasic solution is stirred at room temperature for 6 hours. A solution of FMOC-Cl (11 mmol) in THF is then added dropwise at room temperature over 15 minutes. After 1 hr the solution is acidified with 10% HCl to pH 2, and extracted with EtOAc, washed with water, brine and dried with MgSO₄. The product is purified by flash chromatography.

EXAMPLE 158**2-O-DMT-6-hydroxy-9-FMOC-9-azabicyclo[3.3.1]nonane**

The diol from the previous example (10 mmol) is dissolved in CH₂Cl₂ and pyridine (9:1), cooled to 0 °C and DMT-Cl (10 mmol) is added in portions (1.0 eq). The solution is stirred overnight at 0 °C. The reaction is quenched with MeOH, evaporated, excess solvent removed by azeotropic distillation with toluene. The residue is redissolved in EtOAc, washed with 5% citric acid, NaHCO₃, brine and dried with MgSO₄. The product is purified by flash chromatography on a gradient of MeOH in CH₂Cl₂. The product is obtained in 50% yield, along with 25% ditritylated material which is conserved and recycled, and 25% unreacted starting material.

EXAMPLE 159**25 Dimethyl 3-O-TBDMS-glutarate**

Dimethyl 3-hydroxyglutarate (34 mmol) and imidazole (85 mmol) are dissolved in 150 ml DMF, and TBDMS-Cl added (40 mmol). The solution is stirred overnight, quenched with water (400 ml), extracted with EtOAc, washed with NaHCO₃, brine and dried with MgSO₄.

EXAMPLE 160

- 137 -

3-O-TBDMS-1,5-pentanediol

The material from the previous example is dissolved in dry THF and 35 ml 2 M LiBH₄ in THF added. The solution is heated at reflux for 1.5 hrs, cooled to room temperature and quenched carefully with 10% HCl to pH 2, neutralized with NaHCO₃ and evaporated. The residue is extracted with EtOAc, washed with NaHCO₃, brine and dried with MgSO₄.

EXAMPLE 161**N-Methyl-3-O-TBDMS-7-keto-9-azabicyclo[3.3.1]nonane**

To a stirred solution of oxalyl chloride (10 mmol) in 20 ml THF at -78 °C is added 11 mmol dimethylsulfoxide. The solution is warmed to -35 °C for 5 minutes and re-cooled to -78 °C. A solution of 3-O-TBDMS-1,5-pentanediol (5 mmol) in 10 ml THF is added, and the solution warmed to 0 °C. After 15 minutes, triethylamine (40 mmol) is added and the solution stirred at room temperature to effect complete conversion to the dialdehyde. The solvent is evaporated under reduced pressure, and the crude dialdehyde is redissolved in 40 ml citrate/phosphate buffer (pH 5.5) to which is added 15 mmol methylamine hydrochloride in 50 ml water, followed by the dropwise addition of 12 mmol acetone dicarboxylate in 80 ml water. The solution is stirred 24 hrs, 5 ml of concentrated Hcl is added, and the solution is heated to 90 °C for 1 hr to complete the decarboxylation. Sodium hydroxide is added to pH 12, and the solution extracted with CH₂Cl₂, dried over MgSO₄ and purified by chromatography on alumina.

EXAMPLE 162**N-FMOC-3-O-TBDMS-7-keto-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) is dissolved in 100 ml 1,2-dichloroethane, cooled to 0 °C and 1-chloroethyl chloroformate (22 mmol) is added. The solution is warmed to reflux temperature after 15 minutes, and heated until the starting material is consumed. The solvent is then

- 138 -

removed, and the residue redissolved in anhydrous methanol, and warmed to 50 °C for 1 hour. The solvent is removed and the residue redissolved in 25 ml THF and 25 ml saturated NaHCO₃. A solution of Fmoc-Cl (11 mmol) in THF is then added dropwise at room temperature over 15 minutes. After 1 hr the solution is acidified with 10% HCl to pH 2, and extracted with EtOAc, washed with water, brine and dried with MgSO₄. The product is purified by flash chromatography.

EXAMPLE 163**10 N-Fmoc-3-O-TBDMS-7-hydroxy-9-azabicyclo[3.3.1]nonane**

The material from the previous example (10 mmol) is dissolved in dry THF and 6 ml 2 M LiBH₄ in THF is added. The solution is stirred at room temperature and quenched carefully with 10% HCl to pH 2, neutralized with NaHCO₃ and evaporated.

15 The residue is extracted with EtOAc, washed with NaHCO₃, brine and dried with MgSO₄.

EXAMPLE 164**N-Fmoc-3-O-TBDMS-7-O-DMT-9-azabicyclo[3.3.1]nonane**

The alcohol from the previous example (10 mmol) is dissolved in CH₂Cl₂ and pyridine (9:1), cooled to 0 °C and DMT-Cl (11 mmol) added in portions (1.0 eq). The solution is stirred overnight at room temperature. The reaction is quenched with MeOH, evaporated, and excess solvent removed by azeotropic distillation with toluene. The residue is redissolved in EtOAc, washed with 5% citric acid, NaHCO₃, brine and dried with MgSO₄. The product is purified by silica gael flash chromatography with a gradient of MeOH in CH₂Cl₂.

EXAMPLE 165**N-Fmoc-3-hydroxy-7-O-DMT-9-azabicyclo[3.3.1]nonane**

30 The product of the previous example is added to a solution of triethylamine trihydrofluoride (100 mmol) and triethylamine (50 mmol) in 50 ml THF. The solution is stirred

- 139 -

until the starting material is completely consumed. The reaction is stopped by the addition of saturated NaHCO_3 , followed by evaporation of the solvent. The residue is extracted with EtOAc, washed with NaHCO_3 , brine and dried with MgSO_4 . The product is purified by flash chromatography.

EXAMPLE 166**Synthesis of a trimer containing different aminodiol backbone units**

A shaker flask is charged with 10 μmol (ca. 67 mg) of N-FMOC-2-O-DMT-diethylamino-2'-O-succinylamino TentaGel (abbreviated DMT-FMOC-dea-TG), prepared by succinylation of N-FMOC-2'-hydroxyethyl-2-O-DMT-ethylamine of Example 98, followed by coupling to amino Tentagel using the General Procedure of Example 124. The DMT-FMOC-diethanolamine-TG is treated with 10% piperidine in DMF to remove the FMOC group, washed with DMF, and then with CH_2Cl_2 . The free amine is acylated with cyclohexane carboxylic acid using HATU as the coupling agent following General Procedure of Example 125 and Example 126 to give DMT-de(CH)-TG. The resin is washed with DMF and CH_2Cl_2 , the trityl group removed with 3% TCA in CH_2Cl_2 , and the resin washed with CH_2Cl_2 and pyridine/ CH_3CN (1:1). The resin is then treated with the second scaffold H-phosphonate of Example 81 (abbr. DMT-FMOC-pd-Hphos) and adamantoyl chloride in pyridine/ CH_3CN using the method of the General Procedures of Example 69 and Example 73. After washing with pyridine/ CH_3CN , the resin is treated with a 10% solution of isopropoxyethylamine in carbon tetrachloride/pyridine (1:1) for 20 minutes. The amine solution is removed and the resin washed with DMF, followed by 10% piperidine in DMF to remove the second FMOC group. After washing the resin with DMF, the dimer is acylated with Thiophene carboxylic acid and HATU using the General Procedure. After the acylation is complete, the resin is again washed with DMF, CH_2Cl_2 and detritylated

- 140 -

with 3% TCA in CH_2Cl_2 , and then washed with pyridine/ CH_3CN (1:1). The resin is then treated with the third scaffold H-phosphonate of Example 59 (abbr. DMT-FMOC-hp-Hphos) and adamantoyl chloride in pyridine/ CH_3CN using the method of the General Procedures of Example 69 and Example 73. After washing with pyridine/ CH_3CN , the resin is treated with a 10% solution of 3-aminomethylpyridine in carbon tetrachloride/pyridine (1:1) for 20 minutes. The amine solution is removed and the resin washed with DMF, followed by 10% piperidine in DMF to remove the third FMOC group. After washing the resin with DMF, the trimer is acylated with 4-nitrobenzoic acid and HATU using the General Procedure. The resin is washed with DMF, CH_2Cl_2 , and detritylated with 3% TCA in CH_2Cl_2 . The resin is washed with CH_2Cl_2 to remove traces of acid, and the solid support treated with NH_4OH to cleave the product from the solid support. The ammonia solution is removed in vacuo to give the final compound composed of three different scaffolds (1: diethanolamine, 2: piperidinediol and 3: hydroxyprolinol) acylated with three different carboxylic acids (1: cyclohexane carboxylic acid, 2: thiophene carboxylic acid, and 3: 4-nitrobenzoic acid) and linked with two different substituted phosphoramidates (1-2: isopropoxyethylamine, and 2-3: 3-aminomethylpyridine).

EVALUATION

PROCEDURE 1

Antimicrobial Assay

Staphylococcus aureus

Staphylococcus aureus is known to cause localized skin infections as a result of poor hygiene, minor trauma, psoriasis or eczema. It also causes respiratory infections, pneumonia, toxic shock syndrome and septicemia. It is a common cause of acute food poisoning. It exhibits rapid emergence of drug resistance to penicillin, cephalosporin, vancomycin and nafcillin.

- 141 -

In this assay, the strain *S. aureus* ATCC 25923 (American Type Culture Collection) is used in the bioassay. To initiate the exponential phase of bacterial growth prior to the assay, a sample of bacteria grown overnight at 37 °C in typtocase soy broth (BBL) is used to reinoculate sample wells of 96-well microtiter plates. The assays are carried out in the 96-well microtiter plates in 150 µL volume with approximately 1×10^6 cells per well.

Bacteria in typtocase soy broth (75 µL) is added to the compound mixtures of the invention in solution in 75 µ water in the individual well of the microtiter plate. Final concentrations of the compound mixtures are 25 µM, 10 µM and 1µM. Each concentration of the compound mixtures is assayed in triplicate. The plates are incubated at 37 °C and growth monitored over a 24 hour period by measuring the optical density at 595 nm using a BioRad model 3550 UV microplate reader. The percentage of growth relative to a well containing no compound is determined. Ampicillin and tetracycline antibiotic positive controls are concurrently tested in each screening assay.

PROCEDURE 2

Antimicrobial Mechanistic Assay

Bacterial DNA Gyrase

DNA gyrase is a bacterial enzyme which can introduce negative supercoils into DNA utilizing the free energy derived from ATP hydrolysis. This activity is critical during DNA replication and is a well characterized target for antibiotic inhibition of bacterial growth. In this assay, libraries of compounds of the invention are screened for inhibition of DNA gyrase. The assay measures the supercoiling of a relaxed plasmid by DNA gyrase as an electrophoretic shift on an agarose gel. Initially all library pools are screened for inhibitory activity at 30 µM

- 142 -

and then a dose response analysis is effected with active subsets. Novobiocin, an antibiotic that binds to the β subunit of DNA gyrase is used as a positive control in the assay. The sensitivity of the DNA gyrase assay was determined by titrating the concentration of the known DNA gyrase inhibitor, Novobiocin, in the supercoiling assay. The IC_{50} was determined to be 8 nM, sufficient to identify the activity of a single active species of comparable activity in a library having 30 μ M concentration.

10 PROCEDURE 3

Use of a combinatorial library for identifying of metal chelators and imaging agents

This procedure is used to identify specific compounds of the invention contained in libraries of compounds which each contain a ring which contains an ultraviolet chromophore. The diversity groups attached to the compound bridge are selected from metal binders, coordinating groups such as amine, hydroxyl and carbonyl groups, and other groups having lone pairs of electrons, such that the oligomeric compounds of the invention can form coordination complexes with heavy metals and imaging agents. The procedure is used to identify oligomeric compounds of the invention for chelating and removing heavy metals from industrial broths, waste stream eluents, heavy metal poisoning of farm animals and other sources of contaminating heavy metals, and for use in identifying imaging agent carriers, such as carriers for technetium 99.

An aliquot of a test solution having the desired ion or imaging agent at a known concentration is added to an aliquot of standard solution of the pool of compounds of the invention being assayed. The UV spectrum of this aliquot is measured and is compared to the UV spectrum of a further aliquot of the same solution lacking the test ion or imaging

- 143 -

agent. A shift in the extinction coefficient is indicative of binding of the metal ion or imaging ion to a compound in the library pool being assayed.

PROCEDURE 4

5 Assay of combinatorial library for PLA₂ inhibitors

A preferred target for assay of combinatorially generated pools of compounds is the phospholipase A₂ family. Phospholipases A₂ (PLA₂) are a family of enzymes that hydrolyze the sn-2 ester linkage of membrane phospholipids resulting in release of a free fatty acid and a lysophospholipid (Dennis, E.A., *The Enzymes*, Vol. 16, pp. 307-353, Boyer, P. D., ed., Academic Press, New York, 1983). Elevated levels of type II PLA₂ are correlated with a number of human inflammatory diseases. The PLA₂-catalyzed reaction is the rate-limiting step in the release of a number of pro-inflammatory mediators. Arachidonic acid, a fatty acid commonly linked at the sn-2 position, serves as a precursor to leukotrienes, prostaglandins, lipoxins and thromboxanes. The lysophospholipid can be a precursor to platelet-activating factor. PLA₂ is regulated by pro-inflammatory cytokines and, thus, occupies a central position in the inflammatory cascade (Dennis, *ibid.*; Glaser et al., *TiPs Reviews* 1992, 14, 92; and Pruzanski et al., *Inflammation* 1992, 16, 451). All mammalian tissues evaluated thus far have exhibited PLA₂ activity. At least three different types of PLA₂ are found in humans: pancreatic (type I), synovial fluid (type II) and cytosolic. Studies suggest that additional isoenzymes exist. Type I and type II, the secreted forms of PLA₂, share strong similarity with phospholipases isolated from the venom of snakes. The PLA₂ enzymes are important for normal functions including digestion, cellular membrane remodeling and repair, and in mediation of the inflammatory response. Both cytosolic and

- 144 -

type II enzymes are of interest as therapeutic targets. Increased levels of the type II PLA₂ are correlated with a variety of inflammatory disorders including rheumatoid arthritis, osteoarthritis, inflammatory bowel disease and septic shock, suggesting that inhibitors of this enzyme would have therapeutic utility. Additional support for a role of PLA₂ in promoting the pathophysiology observed in certain chronic inflammatory disorders was the observation that injection of type II PLA₂ into the footpad of rats (Vishwanath et al., *Inflammation* 1988, 12, 549) or into the articular space of rabbits (Bomalaski et al., *J. Immunol.* 1991, 146, 3904) produced an inflammatory response. When the protein was denatured before injection, no inflammatory response was produced.

The type II PLA₂ enzyme from synovial fluid is a relatively small molecule (about 14 kD) and can be distinguished from type I enzymes (e.g. pancreatic) by the sequence and pattern of its disulfide bonds. Both types of enzymes require calcium for activity. The crystal structures of secreted PLA₂ enzymes from venom and pancreatic PLA₂, with and without inhibitors, have been reported (Scott et al., *Science* 1990, 250, 1541). Recently, the crystal structure of PLA₂ from human synovial fluid has been determined (Wery et al., *Nature* 1991, 352, 79). The structure clarifies the role of calcium and amino acid residues in catalysis. Calcium acts as a Lewis acid to activate the scissile ester carbonyl bond of 1,2-diacylglycerophospholipids and binds to the lipid, and a His-Asp side chain dyad acts as a general base catalyst to activate a water molecule nucleophile. This is consistent with the absence of any acyl enzyme intermediates, and is also comparable to the catalytic mechanism of serine proteases. The catalytic residues and the calcium ion are at the end of a deep cleft (ca. 14 Å) in the enzyme. The walls of this cleft contact the hydrocarbon portion of the

phospholipid and are composed of hydrophobic and aromatic residues. The positively-charged amino-terminal helix is situated above the opening of the hydrophobic cleft. Several lines of evidence suggest that the N-terminal portion is the interfacial binding site (Achari et al., Cold Spring Harbor Symp. Quant. Biol. 1987, 52, 441; Cho et al., J. Biol. Chem. 1988, 263, 11237; Yang et al., Biochem. J. 1989, 262, 855; and Noel et al., J. Am. Chem. Soc. 1990, 112, 3704).

Much work has been reported in recent years on the study of the mechanism and properties of PLA₂-catalyzed hydrolysis of phospholipids. In *in vitro* assays, PLA₂ displays a lag phase during which the enzyme adsorbs to the substrate bilayer and a process called interfacial activation occurs. This activation may involve desolvation of the enzyme/lipid interface or a change in the physical state of the lipid around the cleft opening. Evidence favoring this hypothesis comes from studies revealing that rapid changes in PLA₂ activity occur concurrently with changes in the fluorescence of a membrane probe (Burack et al., Biochemistry 1993, 32, 583). This suggests that lipid rearrangement is occurring during the interfacial activation process. PLA₂ activity is maximal around the melting temperature of the lipid, where regions of gel and liquid-crystalline lipid coexist. This is also consistent with the sensitivity of PLA₂ activity to temperature and to the composition of the substrate, both of which can lead to structurally distinct lipid arrangements separated by a boundary region. Fluorescence microscopy was used to simultaneously identify the physical state of the lipid and the position of the enzyme during catalysis (Grainger et al., FEBS Lett. 1989, 252, 73). These studies clearly show that PLA₂ binds exclusively at the boundary region between liquid and solid phase lipid. While the hydrolysis of the secondary ester bond of 1,2-diacylglycerophospholipids catalyzed by the enzyme is rela-

- 146 -

tively simple, the mechanistic and kinetic picture is clouded by the complexity of the enzyme-substrate interaction. A remarkable characteristic of PLA₂ is that maximal catalytic activity is observed on substrate that is aggregated (i.e. phospholipid above its critical micelle concentration), while low levels of activity are observed on monomeric substrate. As a result, competitive inhibitors of PLA₂ either have a high affinity for the active site of the enzyme before it binds to the substrate bilayer or partition into the membrane and compete for the active site with the phospholipid substrate. Although a number of inhibitors appear to show promising inhibition of PLA₂ in biochemical assays (Yuan et al., *J. Am. Chem. Soc.* **1987**, 109, 8071; Lombardo et al., *J. Biol. Chem.* **1985**, 260, 7234; Washburn et al., *J. Biol. Chem.* **1991**, 266, 5042; Campbell et al., *J. Chem. Soc., Chem. Commun.* **1988**, 1560; and Davidson et al., *Biochem. Biophys. Res. Commun.* **1986**, 137, 587), reports describing in vivo activity are limited (Miyake et al., *J. Pharmacol. Exp. Ther.* **1992**, 263, 1302).

In one preferred embodiment, oligomeric compounds of the invention are selected for their potential to interact with, and preferably inhibit, the enzyme PLA₂. Thus, compounds of the invention can be used for topical and/or systemic treatment of inflammatory diseases including atopic dermatitis and inflammatory bowel disease. In selecting the functional groups, advantage can be taken of PLA₂'s preference for anionic vesicles over zwitterionic vesicles. Preferred compounds of the invention for assay for PLA₂ include those having aromatic diversity groups to facilitate binding to the cleft of the PLA₂ enzyme (Oinuma et al., *J. Med. Chem.* **1991**, 34, 2260; Marki et al., *Agents Actions* **1993**, 38, 202; and Tanaka et al., *J. Antibiotics* **1992**, 45, 1071). Benzyl and 4-hexylbenzyl groups are preferred aromatic diversity groups. PLA₂-directed oligomeric compounds of the invention can

- 147 -

further include hydrophobic functional groups such as tetraethylene glycol groups. Since the PLA₂ enzyme has a hydrophobic channel, hydrophobicity is believed to be an important property of inhibitors of the enzyme.

5 After each round of synthesis as described in the above examples, the resulting pools of compounds are screened for inhibition of human type II PLA₂ enzymatic activity. The assay is effected at the conclusion of synthesis to identify the winning compounds of that synthesis. Concurrently, the
10 libraries additionally can be screened in other *in vitro* assays to determine further mechanisms of inhibition.

The pools of the oligomeric compound libraries are screened for inhibition of PLA₂ in the assay using *E. coli* labeled with ³H-oleic acid (Franson et al., *J. Lipid Res.*
15 1974, 15, 380; and Davidson et al., *J. Biol. Chem.* 1987, 262, 1698) as the substrate. Type II PLA₂ (originally isolated from synovial fluid), expressed in a baculovirus system and partially purified, serves as a source of the enzyme. A series of dilutions of each the library pools is done in
20 water: 10 µl of each pool is incubated for 5 minutes at room temperature with a mixture of 10 µl PLA₂, 20 µl 5X PLA₂ Buffer (500 mM Tris 7.0-7.5, 5 mM CaCl₂), and 50 µl water. Samples of each pool are run in duplicate. At this point, 10 µl of ³H
25 *E. coli* cells is added. This mixture is incubated at 37°C for 15 minutes. The enzymatic reaction is stopped with the addition of 50 µl 2M HCl and 50 µl fatty-acid-free BSA (20 mg/ml PBS), vortexed for 5 seconds, and centrifuged at high speed for 5 minutes. 165 µl of each supernate is then put
30 into a scintillation vial containing 6 ml of scintillant (ScintiVerse) and cpms are measured in a Beckman Liquid Scintillation Counter. As a control, a reaction without the combinatorial pool is run alongside the other reactions as well as a baseline reaction containing no oligomeric compounds as well as no PLA₂ enzyme. CPMs are corrected for
35 by subtracting the baseline from each reaction data point.

- 148 -

Confirmation of the "winners" is made to confirm that the oligomeric compound binds to enzyme rather than substrate and that the inhibition of any oligomeric compound selected is specific for type II PLA₂. An assay using ¹⁴C-phosphatidyl ethanolamine (¹⁴C-PE) as substrate, rather than *E. coli* membrane, is used to insure enzyme rather than substrate specificity. Micelles of ¹⁴C-PE and deoxycholate are incubated with the enzyme and oligomer. ¹⁴C-labeled arachidonic acid released as a result of PLA₂-catalyzed hydrolysis is separated from substrate by thin layer chromatography and the radioactive product is quantitated. The "winner" is compared to phosphatidyl ethanolamine, the preferred substrate of human type II PLA₂, to confirm its activity. PLA₂ from other sources (snake venom, pancreatic, bee venom) and phospholipase C, phospholipase D and lysophospholipase can be used to further confirm that the inhibition is specific for human type II PLA₂.

PROCEDURE 5

Probes for the detection of specific proteins and mRNA in biological samples

For the reliable, rapid, simultaneous quantification of multiple varieties of proteins or mRNA in a biological sample without the need to purify the protein or mRNA from other cellular components, a protein or mRNA of interest from a suitable biological sample, i.e., a blood borne virus, a bacterial pathogen product in stool, urine and other like biological samples, is identified using standard microbiological techniques. A probe comprising an oligomeric compound of the invention is identified by a combinatorial search as noted in the above examples. Preferred for the mRNA probe are compounds synthesized to include "nucleobase" diversity groups (adenine, guanine, thymine and cytosine as the letters) complementary to at least a portion of the

- 149 -

nucleic acid sequence of the mRNA. Preferred for the protein probe are compounds synthesized to include chemical functional groups that act as hydrogen bond donors and acceptors, sulfhydryl groups, hydrophobic lipophilic moieties capable of hydrophobic interactions groups and groups capable of ionic interactions. The probe is immobilized on insoluble CPG solid support utilizing the procedure of Pon, R. T., Protocols for Oligonucleotides and Analogs, Agrawal, S., Ed., Humana Press, Totowa, NJ, 1993, p 465-496. A known aliquot of the biological sample under investigation is incubated with the insoluble CPG support having the probe thereon for a time sufficient to hybridize the protein or mRNA to probe and thus to link them via the probe to the solid support. This immobilizes protein or mRNA present in the sample to the CPG support. Other non-immobilized materials and components are then washed off the CPG with a wash media suitable for use with the biological sample. The mRNA on the support is labeled with ethidium bromide, biotin or a commercial radionucleotide and the amount of label immobilized on the CPG support is measured to indicate the amount of mRNA present in the biological sample. In a similar a protein is also labeled and quantified.

PROCEDURE 6

Leukotriene B₄ assay

Leukotriene B₄ (LTB₄) has been implicated in a variety of human inflammatory diseases, and its pharmacological effects are mediated via its interaction with specific surface cell receptors. Library subsets are screened for competitive inhibition of radiolabeled LTB₄ binding to a receptor preparation.

A NenquestTM Drug Discovery System Kit (NEN Research Products, Boston, MA) is used to select an inhibitor of the interaction of Leukotriene B₄ (LTB₄) with receptors on a preparation of guinea pig spleen membrane. [³H] Leukotriene

- 150 -

B₄ reagent is prepared by adding 5 mL of ligand diluent (phosphate buffer containing NaCl, MgCl₂, EDTA and Bacitracin, pH 7.2) to 0.25 mL of the radioligand. The receptor preparation is made by thawing the concentrate, adding 35 mL of ligand diluent and swirling gently in order to resuspend the receptor homogeneously. Reagents are kept on ice during the course of the experiment, and the remaining portions are stored at -20°C.

Library subsets prepared as per general procedure of examples above are diluted to 5 µM, 50 µM and 500 µM in phosphate buffer (1x PBS, 0.1% azide and 0.1% BSA, pH 7.2), yielding final test concentrations of 0.5 µM, 5 µM and 50 µM, respectively. Samples are assayed in duplicate. [³H] LTB₄ (25 µL) is added to 25 µL of either appropriately diluted standard (unlabeled LTB₄) or library subset. The receptor suspension (0.2 mL) is added to each tube. Samples are incubated at 4°C for 2 hours. Controls include [³H] LTB₄ without receptor suspension (total count vials), and sample of ligand and receptor without library molecules (standard).

After the incubation period, the samples are filtered through GF/B paper that had been previously rinsed with cold saline. The contents of each tube are aspirated onto the filter paper to remove unbound ligand from the membrane preparation, and the tubes washed (2 x 4 mL) with cold saline. The filter paper is removed from the filtration unit and the filter disks are placed in appropriate vials for scintillation counting. Fluor is added, and the vials shaken and allowed to stand at room temperature for 2 to 3 hours prior to counting. The counts/minute (cpm) obtained for each sample are subtracted from those obtained from the total count vials to determine the net cpm for each sample. The degree of inhibition of binding for each library subset is determined relative to the standard (sample of ligand and receptor without library molecules).

Each of the published documents mentioned in this


- 151 -

specification are herein incorporated in their entirety.

Those skilled in the art will appreciate that numerous changes and modifications may be made to the preferred embodiments of the invention and that such changes and modifications may be made without departing from the spirit of the invention. It is therefore intended that the appended claims cover all such equivalent variations as fall within the true spirit and scope of the invention.

WHAT IS CLAIMED IS:

1. An oligomeric compound comprising a plurality of aminodiol monomer subunits joined by linking groups, wherein each of said aminodiol monomer subunits has one of the structures I, II, III, IV, V, VI, VII, VIII, IX, X, or XI;




wherein:

each x is, independently, 0 to 5;

na , nb and nc are each, independently, 0 to 2, where the sum of na , nb and nc is from 1 to 5;

R_1 is -T-L or a base labile protecting group;

T is a single bond, a methylene group or a group having formula:



wherein:


- 5 R_{10} is =O, =S, or =NR₁₁;
 R_5 and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;
each R₆, R₇, R₈, R₉, R₁₁, R₁₂ and R₁₃ are,
independently, H, alkyl or haloalkyl having 1 to about 10
10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;
m and n, independently, are 0 to 5;
p is 0 or 1;
15 q is 1 to about 10;
L is H, substituted or unsubstituted C₂-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted C₄-C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇
20 carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a
25 metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂),

- 154 -

sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups;

R_2 is hydrogen or C_1 - C_{10} alkyl;

R_3 and R_4 are independently hydrogen, an acid labile hydroxyl protecting group, a linking group or a conjugate group, wherein said linking group has the formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_0)T_0$;

Y_0 is H or $[Q_2]_j-Z_2$;

T_0 is $[Q_1]_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_4 - C_7 carbocyclo alkyl C_4 - C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7 - C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H, C_1 - C_2 alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, C_6 - C_{14} aryl, C_7 - C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group; and

- 155 -

provided that at least one of said aminodiol monomer subunits in said oligomeric compound does not have structure III.

2. The oligomeric compound of claim 1 wherein said J₁ is =O or =S and said J₂ is OH.

3. The oligomeric compound of claim 1 wherein said J₁ is =O, said J₂ is N(Y₀)T₀ and at least two of said N(Y₀)T₂ are the same.

4. The oligomeric compound of claim 1 wherein said J₁ is =O, said J₂ is N(Y₀)T₀ and wherein at least two of said N(Y₀)T₀ are different.

5. The oligomeric compound of claim 1 wherein each of said R₁ are the same.


6. The oligomeric compound of claim 1 wherein at least two of said R₁ are different.

7. The oligomeric compound of claim 1 wherein each of said aminodiol monomer subunits are the same.

8. The oligomeric compound of claim 1 wherein at least two of said aminodiol monomer subunits are different.

- 156 -

9. A library of oligomers, each of said oligomers comprising a plurality of aminodiols monomer subunits joined by linking groups, said aminodiols monomer subunits, each of said subunits having structure I, II, III, IV, V, VI, VII, VIII, IX, X, or XI;



wherein:


each x is, independently, 0 to 5;

na , nb and nc are each, independently, 0 to 2, where the sum of na , nb and nc is from 1 to 5;

R_1 is -T-L or a base labile protecting group;

- 157 -

T is a single bond, a methylene group or a group having formula:



5 wherein:

R_{10} is =O, =S, or =NR₁₁;

R_5 and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;

each R_6 , R_7 , R_8 , R_9 , R_{11} , R_{12} and R_{13} are,

10 independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

15 p is 0 or 1;

q is 1 to about 10;


L is H, substituted or unsubstituted C₂-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted C₁-C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇ carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂),

- 158 -

sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups;

R_2 is hydrogen or C_1 - C_{10} alkyl;

R_3 and R_4 are independently hydrogen, an acid labile hydroxyl protecting group, a linking group or a conjugate group, wherein said linking group has the formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_0)T_0$;

Y_0 is H or $[Q_2]_j-Z_2$;

T_0 is $[Q_1]_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_4 - C_7 carbocyclo alkyl C_4 - C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7 - C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H, C_1 - C_2 alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, C_6 - C_{14} aryl, C_7 - C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group; and

- 159 -

provided that at least one of said aminodiol monomer subunits in each oligomeric compound of said library does not have structure III.

10. The library of claim 9 wherein said J_1 is =O or =S and said J_2 is OH.

11. The library of claim 9 wherein said J_1 is =O, said J_2 is $N(Y_0)T_0$ and at least two of said $N(Y_0)T_0$ are the same.

12. The library of claim 9 wherein said J_1 is =O, said J_2 is $N(Y_0)T_0$ and at least two of said $N(Y_0)T_0$ are different.


13. The library of claim 9 wherein each of said R_1 is the same.

14. The library of claim 9 wherein at least two of said R_1 are different.

15. A method for preparing an oligomer comprising:

(a) selecting an aminodiol monomer subunit having the structure I , II, III, IV, V, VI, VII, VIII, IX, X, or XI;

- 160 -



wherein:

each x is, independently, 0 to 5;

na , nb and nc are each, independently, 0 to 2, where the sum of na , nb and nc is from 1 to 5;

R_1 is a base labile amino protecting group;

R_2 is hydrogen or $\text{C}_1\text{-C}_{10}$ alkyl;

one of R_3 or R_4 is hydrogen or an activated phosphite group and the other of R_3 or R_4 is an acid labile hydroxyl protecting group;

(b) attaching said aminodiol monomer subunit to a

- 161 -


solid support to form a solid support bound aminodiol monomer subunit;

(c) treating said acid labile hydroxyl protecting group with a dilute acid to form a free hydroxyl group;

(d) reacting said free hydroxyl group with a further aminodiol monomer subunit having structure I, II, III, IV, V, VI, or VII, VIII, IX, X, or XI thereby forming an oligomeric compound bound to said solid support, said oligomeric compound containing a phosphite linkage;

(e) optionally iteratively repeating steps (c) and (d) to increase the length of the oligomeric compound bound to said solid support;

(f) optionally, prior to step (c) or after step (d) oxidizing said phosphite linkage to form a phosphate linking group wherein said linking groups are selected having formula:



wherein:

J_1 is =O or =S;

J_2 is OH or N(Y₀)T₀;

Y₀ is H or [Q₂]_j-Z₂;

T₀ is [Q₁]_k-Z₁, or together Y₀ and T₀ are joined in a nitrogen heterocycle;

Q₁ and Q₂ independently are C₂-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₄-C₇ carbocyclo alkyl C₄-C₇ carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C₇-C₁₄ aralkyl;

j and k independently are 0 or 1;

Z₁ and Z₂ independently are H, C₁-C₂ alkyl, C₂-C₂₀


- 162 -

alkenyl, C₂-C₂₀ alkynyl, C₆-C₁₄ aryl, C₇-C₁₅ aralkyl, halogen, CH=O, OR₁₂, SR₁₂, NR₁₂R₁₃, C(=NH)NR₁₂R₁₃, CH(NR₁₂R₁₃), NHC(=NH)NR₁₂R₁₃, CH(NH₂)C(=O)OH, C(=O)NR₁₂R₁₃, C(=O)OR₁₂, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group;

(g) prior to step (e) or after step (f) contacting said solid support bound aminodiol monomer subunit or said support bound oligomeric compound with a base to remove said base labile amino protecting group to form the solid support bound aminodiol monomer subunit or support bound oligomeric compound having a free amine, and derivatizing said free amine with a group of the formula:

wherein:

T is a single bond, a methylene group or a group having formula:



where:

R₁₀ is =O, =S, or =NR₁₁;

R₅ and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;

each R₆, R₇, R₈, R₉, R₁₁, R₁₂ and R₁₃ are, independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

p is 0 or 1;

q is 1 to about 10;

L is H, substituted or unsubstituted C₂-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted C₄-

- 163 -

C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇ carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups;

(h) optionally repeating steps (c) and (d) followed by step (g) to increase the length of the oligomeric compound bound to said solid support;

(i) treating said oligomeric compound bound to said solid support with acid to deprotect any protecting groups; and

(j) cleaving said oligomeric compound from said solid support.

16. The process of claim 15 wherein said step (g) is conducted after said step (b).

17. The process of claim 15 wherein said step (g) is conducted prior to step (d) for the addition of at least one monomeric subunit to said oligomeric compound.

- 164 -

18. The process of claim 15 wherein said step (g) is conducted prior to each iteration of said step (d).


19. The process of claim 15 wherein said step (g) is conducted only after at least one iteration of said step (e).

20. The process of claim 15 wherein said step (g) is conducted after said step (f) for the addition of at least one monomeric subunit to said oligomeric compound.

21. A method for preparing a combinatorial library comprising:

(a) selecting a plurality of aminodiols monomer subunits having the structure I, II, III, IV, V, VI, VII, VIII, IX, X, or XI:

- 165 -



wherein:

each x is, independently, 0 to 5;

na , nb and nc are each, independently, 0 to 2, where the sum of na , nb and nc is from 1 to 5;

R_1 is a base labile amino protecting group;

R_2 is hydrogen or $\text{C}_1\text{-C}_{10}$ alkyl;

one of R_3 or R_4 is hydrogen or an activated phosphite group and the other of R_3 or R_4 is an acid labile hydroxyl protecting group;

(b) attaching said aminodiol monomer subunits to a

- 166 -


solid support to form a solid support bound aminodiol monomer subunits;

(c) treating said acid labile hydroxyl protecting groups with a dilute acid to form a free hydroxyl groups;

5 (d) reacting said free hydroxyl groups with further aminodiol monomer subunits having structure I, II, III, IV, V, VI, VII, VIII, IX, X, or XI thereby forming an oligomeric compound bound to said solid support, said oligomeric compound containing a phosphite linkage;

10 (e) optionally iteratively repeating steps (c) and (d) to increase the length of the oligomeric compound bound to said solid support;

(f) optionally, prior to step (c) or after step (d) oxidizing said phosphite linkage to form phosphate
15 linking groups having formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_0)T_0$;

20 Y_0 is H or $[Q_2]_j-Z_2$;

T_0 is $[Q_1]_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_4-C_7 carbocyclo alkyl C_4-C_7 carbocyclo
25 alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7-C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H, C_1-C_2 alkyl, C_2-C_{20}
30 alkenyl, C_2-C_{20} alkynyl, C_6-C_{14} aryl, C_7-C_{15} aralkyl, halogen,

- 167 -


CH=O, OR₁₂, SR₁₂, NR₁₂R₁₃, C(=NH)NR₁₂R₁₃, CH(NR₁₂R₁₃),
 NHC(=NH)NR₁₂R₁₃, CH(NH₂)C(=O)OH, C(=O)NR₁₂R₁₃, C(=O)OR₁₂, a metal
 coordination group, a reporter group, a nitrogen-containing
 heterocycle, a purine, a pyrimidine, a phosphate group, a
 5 polyether group, or a polyethylene glycol group;

(g) prior to step (e) or after step (f) contacting
 said solid support bound aminodiol monomer subunits or said
 support bound oligomeric compounds with a base to remove said
 base labile amino protecting groups to form the solid support
 10 bound aminodiol monomer subunits or support bound oligomeric
 compounds having a free amine, and derivatizing said free
 amine with a group of the formula



wherein:

15 T is a single bond, a methylene group or a group
 having formula:



where:

20 R₁₀ is =O, =S, or =NR₁₁;
 R₅ and E, independently, are a single bond, CH=CH,
 C≡C, O, S, NR₁₁, or C₆-C₁₄ aryl;

each R₆, R₇, R₈, R₉, R₁₁, R₁₂ and R₁₃ are,
 independently, H, alkyl or haloalkyl having 1 to about 10
 25 carbon atoms, alkenyl having 2 to about 10 carbon atoms,
 alkynyl having 2 to about 10 carbon atoms, or aryl having 7
 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

p is 0 or 1;

30 q is 1 to about 10;

L is H, substituted or unsubstituted C₂-C₁₀ alkyl,
 substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or
 unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted C₄-

05973381-032598

- 168 -

- C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇ carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide (SO), sulfone (SO₂), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups;
- (h) optionally repeating steps (c) and (d) followed by step (g) to increase the length of the oligomeric compounds bound to said solid support;
- (h) treating said oligomeric compounds bound to said solid support with acid to deprotect any protecting groups; and
- (i) cleaving said oligomeric compounds from said solid support.

22. The process of claim 21 wherein said step (g) is conducted after said step (b).

23. The process of claim 21 wherein said step (g) is conducted prior to step (e) for the addition of at least one monomeric subunit to each of said oligomeric compounds.

- 169 -

24. The process of claim 21 wherein said step (g) is conducted prior to each iteration of said step (e).

25. The process of claim 21 wherein said step (g) is conducted only after at least one iteration of said step (e).

5 26. The process of claim 21 wherein said step (g) is conducted after said step (f) for the addition of at least one monomeric subunit to each of said oligomeric compounds.

065220-132620

Applicant or Patentee: Normand Hebert

Serial No.: PCT/US96/09604

Attorney's Docket No.: ISIS-2297

Date Filed or Issued: June 7, 1996

For: Combinatorial Libraries Having Aminodiol Monomer Subunits

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(d) and 1.27(c)) - SMALL BUSINESS CONCERN**

I hereby declare that I am:

() the owner of the small business concern identified below:

(XX) an official empowered to act on behalf of the concern
identified below:

NAME OF CONCERN: ISIS PHARMACEUTICALS, INC.

ADDRESS OF CONCERN: 2292 FARADAY AVENUE, CARLSBAD, CALIFORNIA 92008

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR 121.12, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, in that: (1) the number of employees of the concern, including those of its affiliates, does not exceed 500 persons; and (2) the concern has not assigned, granted, conveyed, or licensed, and is under no obligation under contract or law to assign, grant, convey, or license, any rights in the invention to any person who could not be classified as an independent inventor if that person had made the invention, or to any concern which would not qualify as a small business concern or a nonprofit organization under this section. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention entitled **Combinatorial Libraries Having Aminodiol Monomer Subunits** by inventor(s) **Normand Hebert** described in

() specification filed herewith.

(XX) international application serial no. PCT/US96/09604, filed June 7, 1996.

If the rights held by the above-identified small business concern are not exclusive, each individual, concern or organization having rights in the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

***NOTE:** Separate verified statements are required for each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

FULL NAME:

ADDRESS:

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

FULL NAME:

ADDRESS:

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING:
TITLE OF PERSON SIGNING:
ADDRESS OF PERSON SIGNING:


B. Lynne Parshall
Executive Vice President
2292 Faraday Avenue,
Carlsbad, California 92008


SIGNATURE


November 25, 1997
DATE

1/2

Fig. 1A



2/2

Fig. 1B

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and

I verily believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **COMBINATORIAL LIBRARIES HAVING AMINODIOL MONOMER SUBUNITS**, the specification of which:

() is attached hereto.

(XX) was filed on June 7, 1996 as International Application Serial No. PCT/US96/09604.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to the patentability of this application in accordance with 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of any application on which priority is claimed:

Country	Number	Date Filed	Priority Claimed
_____	_____	_____	_____
_____	_____	_____	_____
_____	_____	_____	_____

PTO/PCT Rec'd 25 MAR 1996

DOCKET NO. ISIS-2297

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

Normand Hebert

U.S. Serial No.: 08/973,381

International Serial No.: PCT/US96/09604

Filed: June 7, 1996

For: Combinatorial Libraries Having Aminodiol
Monomer Subunits

Assistant Commissioner for Patents
Washington DC 20231

Sir:

ASSOCIATE POWER OF ATTORNEY


The undersigned, of the firm WOODCOCK WASHBURN KURTZ
MACKIEWICZ & NORRIS LLP, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania
19103, Attorney and/or Agents for Applicant(s), hereby appoints the following:

Robert B. Washburn	Registration No. <u>16,574</u>	Michael P. Dunnam	Registration No. <u>32,611</u>
Richard E. Kurtz	Registration No. <u>19,263</u>	Michael D. Stein	Registration No. <u>34,734</u>
John J. Mackiewicz	Registration No. <u>19,709</u>	Albert J. Marcellino	Registration No. <u>34,664</u>
Norman L. Norris	Registration No. <u>24,196</u>	David R. Bailey	Registration No. <u>35,057</u>
Albert W. Preston, Jr.	Registration No. <u>25,366</u>	Barbara L. Mullin	Registration No. <u>38,250</u>
Dale M. Heist	Registration No. <u>28,425</u>	Lynn A. Malinoski	Registration No. <u>38,788</u>
Philip S. Johnson	Registration No. <u>27,200</u>	Lori Y. Beardell	Registration No. <u>34,293</u>
John W. Caldwell	Registration No. <u>28,937</u>	Doreen Yatko Trujillo	Registration No. <u>35,719</u>
Gary H. Levin	Registration No. <u>28,734</u>	Paul K. Legaard	Registration No. <u>38,534</u>
Steven J. Rocci	Registration No. <u>30,489</u>	Kevin M. Flannery	Registration No. <u>35,871</u>
Dianne B. Elderkin	Registration No. <u>28,598</u>	David A. Cherry	Registration No. <u>35,099</u>
Francis A. Paintin	Registration No. <u>19,386</u>	Anthony J. Rossi	Registration No. <u>24,053</u>
John P. Donohue, Jr.	Registration No. <u>29,916</u>	Michael J. Swope	Registration No. <u>38,041</u>
Henrik D. Parker	Registration No. <u>31,863</u>	Michael J. Bonella	Registration No. <u>P41,628</u>
Suzanne E. Miller	Registration No. <u>32,279</u>	Harold H. Fullmer	Registration No. <u>P42,560</u>
Lynn B. Morreale	Registration No. <u>32,842</u>	John E. McGlynn	Registration No. <u>P42,863</u>
Mark DeLuca	Registration No. <u>33,229</u>	Gail Ann Dalickas	Registration No. <u>40,979</u>
Joseph Lucci	Registration No. <u>33,307</u>	Kimberly R. Hild	Registration No. <u>39,224</u>

Lawrence A. Aaronson	Registration No. 38,369
Jonathan M. Waldman	Registration No. <u>40,861</u>
Joseph R. Condo	Registration No. <u>P42,431</u>
Michael K. Jones	Registration No. <u>P41,100</u>

his/her associates with full power to prosecute the above-identified application and to transact all business in the Patent Office connected therewith and requests that correspondence continue to be directed to the firm of WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP at the above address.

Date: March 25, 1998


Michael P. Straher
Registration No. 38,325

Woodcock Washburn Kurtz
Mackiewicz & Norris LLP
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone: (215) 568-3100
Facsimile: (215) 568-3439

© 1997 WWKMN

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status (patented, pending)
<u>08/483,311</u>	<u>June 7, 1995</u>	<u>Pending</u>
<u>PCT/US95/00356</u> (U.S. serial no. 08/669,505)	<u>January 11, 1995</u>	<u>Allowed</u>
<u>08/179,970</u>	<u>January 11, 1994</u>	<u>Pending</u>
<u>08/180,134</u>	<u>January 11, 1994</u>	<u>Patented</u>

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: **John W. Caldwell and Michael P. Straher**, Registration Nos. 28,937 and 38,325 of the firm of **WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP**, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania 19103, and **Herb Boswell, Laurel Bernstein and Andrew E. Granston**, Registration Nos. 27,311, 37,280 and 38,473, of **ISIS Pharmaceuticals, Inc.**, 2292 Faraday Avenue, Carlsbad, California 92008.

Address all telephone calls and correspondence to:

Michael P. Straher
WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP
One Liberty Place - 46th Floor
Philadelphia, PA 19103
Telephone No. 215-568-3100.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the

United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1	Full Name: <u>Normand Hebert</u>	Inventor's Signature: <u>Normand Hebert</u>	Date: <u>12/4/97</u>
	Residence: <u>Cardiff, California</u>	Citizenship: Canada	
	Post Office Address: 1861 Montgomery Avenue Cardiff, California 92007		
2	Full Name:	Inventor's Signature:	Date
	Residence:	Citizenship:	
	Post Office Address:		
3	Full Name:	Inventor's Signature:	Date
	Residence:	Citizenship:	
	Post Office Address:		
4	Full Name:	Inventor's Signature:	Date
	Residence:	Citizenship:	
	Post Office Address:		
5	Full Name:	Inventor's Signature:	Date
	Residence:	Citizenship:	
	Post Office Address:		